

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM15-0044555 | | |
| Date Assigned: | 03/12/2015 | Date of Injury: | 10/22/2012 |
| Decision Date: | 04/22/2015 | UR Denial Date: | 02/20/2015 |
| Priority: | Standard | Application Received: | 03/09/2015 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 35 year old, male patient, who sustained an industrial injury on 10/22/2012. A primary treating office visit dated 02/13/2015, reported chief complaint of lower back pain and left leg sciatica. The patient states that he has chronic pain to the low back that radiates down the left leg all the way to the heel with occasional numbness. He continues to use Norco for pain relief. Prior treatment includes epidural steroid injections, physical therapy and anti-inflammatory medication. Of note, the patient did get surgical recommendation for a total disc replacement at L5-S1. Objective findings showed his lumbar spine has tenderness at the lumbosacral junction as well as in the left lower lumbar spine with mild spasm found. The left sciatic notch is tender. Lumbar flexion brings fingertips to the level of the midtibia and he extends 20 degrees. He reports low back pain as he extends from the flexed over position. He is able to tilt to the right and left 20 degrees with pain reported at each limit. A straight leg raise on the left at 90 degrees reproduces low back and left leg pain. Straight leg raise on the right reproduces low back pain only. The impression noted lumbago with L5-S1 annular tear bilaterally, radiculopathy left greater than right and L5-S1 herniated nucleus pulposus.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MRI of lumbar spine: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints, Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303.

Decision rationale: Regarding the indications for imaging in case of back pain, MTUS guidelines stated: Lumbar spine x rays should not be recommended in patients with low back pain in the absence of red flags for serious spinal pathology, even if the pain has persisted for at least six weeks. However, it may be appropriate when the physician believes it would aid in patient management. Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computer tomography [CT] for bony structures). Furthermore, and according to MTUS guidelines, MRI is the test of choice for patients with prior back surgery, fracture or tumors that may require surgery. The patient does not have any clear evidence of new lumbar nerve root compromise. There is no clear evidence of significant change in the patient signs or symptoms suggestive of new pathology. Therefore, the request for lumbar MRI is not medically necessary.

Norco 5/325mg quantity 120: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints, Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, Norco (Hydrocodone/Acetaminophen) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. According to

the patient file, there is no objective documentation of pain and functional improvement to justify continuous use of Norco. Norco was used for longtime without documentation of functional improvement or evidence of return to work or improvement of activity of daily living. Therefore, the prescription of Norco 5/325mg #120 is not medically necessary.