

Case Number:	CM15-0045975		
Date Assigned:	03/18/2015	Date of Injury:	05/15/2003
Decision Date:	04/23/2015	UR Denial Date:	02/18/2015
Priority:	Standard	Application Received:	03/11/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 55-year-old male who sustained an industrial injury on 05/15/2003. Diagnoses include chronic pain, lumbar radiculitis, post lumbar fusion, and post spinal cord stimulator. Treatment to date has included medications. A physician progress note dated 01/13/2015 documents he has continuing pain in his low back radiating to his left leg, and it is constant and severe in nature. He ambulates with a cane, and range of motion is restricted. There is a mildly positive sitting straight leg raise bilaterally. A hand-written physician progress note dated 02/03/2015 documents the injured worker uses Naproxen for inflammation. The stimulator is working well. His testicles hurt intermittently. He has an antalgic gait and uses a walker. Treatment requested is for Naproxen 500 mg #90.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Naproxen 500 mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS
Page(s): 67, 70-73.

Decision rationale: According to the MTUS Anti-inflammatories are the traditional first line of treatment, to reduce pain so activity and functional restoration can resume, but long-term use may not be warranted. (Van Tulder-Cochrane, 2000) Recommended with cautions below.

Disease-State Warnings for all NSAIDs: All NSAIDs have[U.S. Boxed Warning]: for associated risk of adverse cardiovascular events, including, MI, stroke, and new onset or worsening of pre-existing hypertension. NSAIDs should never be used right before or after a heart surgery (CABG - coronary artery bypass graft). NSAIDs can cause ulcers and bleeding in the stomach and intestines at any time during treatment (FDA Medication Guide). See NSAIDs, GI Symptoms and Cardiovascular Risks. Other disease-related concerns(non-boxed warnings):

Hepatic: Use with caution in injured workers with moderate hepatic impairment and not recommended for injured workers with severe hepatic impairment. Borderline elevations of one or more liver enzymes may occur in up to 15% of injured workers taking NSAIDs. Renal: Use of NSAIDs may compromise renal function. FDA Medication Guide is provided by FDA mandate on all prescriptions dispensed for NSAIDs. Routine Suggested Monitoring: Package inserts for NSAIDs recommend periodic lab monitoring of a CBC and chemistry profile (including liver and renal function tests). There has been a recommendation to measure liver transaminases within 4 to 8 weeks after starting therapy, but the interval of repeating lab tests after this treatment duration has not been established. Routine blood pressure monitoring is recommended.

Overall Dosing Recommendation: It is generally recommended that the lowest effective dose be used for all NSAIDs for the shortest duration of time consistent with the individual injured worker treatment goals. According to the documents available for review, it appears that the injured worker is taking this medication for long-term therapy of a chronic condition. Given the increased risks associated with long-term use of this medication and no documented evidence that the lowest possible dose is being used for the shortest period of time, the requirements for treatment have not been met and medical necessity has not been established.