

<b>Case Number:</b>	CM15-0048393		
<b>Date Assigned:</b>	04/14/2015	<b>Date of Injury:</b>	06/23/2010
<b>Decision Date:</b>	05/11/2015	<b>UR Denial Date:</b>	03/03/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	03/13/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: New York, Tennessee  
 Certification(s)/Specialty: Emergency Medicine

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 44 year old female, who sustained an industrial injury on 06/03/2010. She has reported subsequent left ankle pain and was diagnosed with internal derangement of the left ankle, avulsion fracture of the left ankle at the lateral malleolus, Achilles bursitis or tendinitis and myofascial pain syndrome. Treatment to date has included oral pain medication and physical therapy. In a progress note dated 02/18/2015, the injured worker complained of left ankle pain, swelling, decreased sensation and muscle spasms. Objective findings were notable for global swelling around the entire left ankle, decreased range of motion of the left ankle and muscle spasms and trigger points surrounding the left ankle muscles. A request for authorization of 8 sessions of acupuncture, Flexeril and urine toxicology screen was made.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**8 Acupuncture Sessions:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Acupuncture Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Acupuncture Treatment Guidelines.

**Decision rationale:** Section 9792.24.1 of the California Code of regulations states that Acupuncture is used as an option when pain medication is reduced or not tolerated or as an adjunct to physical rehabilitation. It is the insertion and removal of filiform needles to stimulate acupoints (acupuncture points). Needles may be inserted, manipulated, and retained for a period of time. Acupuncture can be used to reduce pain, reduce inflammation, increase blood flow, increase range of motion, decrease the side effect of medication-induced nausea, promote relaxation in an anxious patient, and reduce muscle spasm. Acupuncture with electrical stimulation is the use of electrical current on the needles at the acupuncture site. It is used to increase effectiveness of the needles by continuous stimulation of the acupoint. Physiological effects (depending on location and settings) can include endorphin release for pain relief, reduction of inflammation, increased blood circulation, analgesia through interruption of pain stimulus, and muscle relaxation. It is indicated to treat chronic pain conditions, radiating pain along a nerve pathway, muscle spasm, inflammation, scar tissue pain, and pain located in multiple sites. Specific indications for treatment of pain include treatment of joint pain, joint stiffness, soft tissue pain and inflammation, paresthesias, post-surgical pain relief, muscle spasm and scar tissue pain. OGD states that acupuncture is not recommended for acute back pain, but is recommended as an option for chronic low back pain in conjunction with other active interventions. Acupuncture is recommended when use as an adjunct to active rehabilitation. Frequency and duration of acupuncture or acupuncture with electrical stimulation may be performed as follows: 1) Time to produce functional improvement: 3 to 6 treatments. 2) Frequency: 1 to 3 times per week. 3) Optimum duration: 1 to 2 months. Acupuncture treatments may be extended if functional improvement is documented. In this case the recommended number of 8 visits surpasses the recommended number of 3 to 6 treatments to produce improvement. The request should not be authorized and is not medically necessary.

**Flexeril 7.5 MG 1 Tab By Mouth TID #Unspecified:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 63.

**Decision rationale:** Flexeril is cyclobenzaprine. Cyclobenzaprine is a muscle relaxant. Cyclobenzaprine is recommended as an option, for a short course of therapy. It has been found to be more effective than placebo with greater adverse side effects. Its greatest effect is in the first 4 days. Treatment should be brief. Non-sedating muscle relaxants are recommended with caution as a second-line option for short-term treatment (less than two weeks) of acute exacerbations in patients with chronic LBP. Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. Sedation is the most commonly reported adverse effect of muscle relaxant medications. These drugs should be used with caution in patients driving motor vehicles or operating heavy machinery. In this case the patient has been

taking Flexeril since at least February 2015. The duration of treatment surpasses the recommended short-term duration of two weeks. The request should not be authorized and is not medically necessary.

**Urine Toxicology Screen:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 78. Decision based on Non-MTUS Citation Official Disability Guidelines: Pain, urine drug testing.

**Decision rationale:** Chronic Pain Medical Treatment Guidelines state that urinary drug testing should be used if there are issues of abuse, addiction, or pain control in patients being treated with opioids. ODG criteria for Urinary Drug testing are recommended for patients with chronic opioid use. Patients at low risk for addiction/aberrant behavior should be tested within 6 months of initiation of therapy and yearly thereafter. Those patients with moderate risk for addiction/aberrant behavior should undergo testing 2-3 times/year. Patients with high risk of addiction/aberrant behavior should be tested as often as once per month. In this case the patient has not been prescribed opioid medication. There is no documentation of aberrant/addictive behavior. There is no medical necessity for urine drug testing at this time. The request should not be authorized and is not medically necessary.