

Case Number:	CM15-0049014		
Date Assigned:	03/20/2015	Date of Injury:	08/01/2001
Decision Date:	05/01/2015	UR Denial Date:	02/27/2015
Priority:	Standard	Application Received:	03/16/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, District of Columbia, Maryland
 Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 46 year old female, who sustained an industrial injury on August 1, 2001. She reported injury of the neck, shoulders, upper extremities, and upper back. The injured worker was diagnosed as having neck sprain, right carpal tunnel syndrome. Treatment to date has included magnetic resonance imaging, medications, electrodiagnostic studies, carpal tunnel release, work modifications, and physical therapy. She was seen on February 4, 2015 by the primary treating physician, for bilateral upper extremity pain, cervical spine, thoracic spine, and lumbar spine pain. She indicates her pain is 4/10 on a pain scale with medications, and 8/10 without medications. The treatment plan includes: continuation of Norco 10/325mg, Methoderm topical cream, Ambien 10mg, and Prilosec 20mg; and recommendations for continuation of medications prescribed as directed by other physicians, continuation of carpal tunnel brace, and follow-up in 2 months or sooner as needed. The request is for Norco 10/325mg, and Methoderm topical cream.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, criteria for use.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 78, 91.

Decision rationale: Per MTUS Chronic Pain Medical Treatment Guidelines p78 regarding on-going management of opioids "Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: Pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug related behaviors. These domains have been summarized as the '4 A's' (Analgesia, activities of daily living, adverse side effects, and any aberrant drug-taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs." Review of the available medical records reveals no documentation to support the medical necessity of norco nor any documentation addressing the '4 A's' domains, which is a recommended practice for the on-going management of opioids. Specifically, the notes do not appropriately review and document pain relief, functional status improvement, appropriate medication use, or side effects. The MTUS considers this list of criteria for initiation and continuation of opioids in the context of efficacy required to substantiate medical necessity, and they do not appear to have been addressed by the treating physician in the documentation available for review. Efforts to rule out aberrant behavior (e.g. CURES report, UDS, opiate agreement) are necessary to assure safe usage and establish medical necessity. UDS dated 11/11/14 was consistent with prescribed medications. As MTUS recommends discontinuing opioids if there is no overall improvement in function, medical necessity cannot be affirmed.

Menthoderm topical cream: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs), Salicylate topicals.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical analgesics Page(s): 60, 105, 111-113.

Decision rationale: Menthoderm is methyl salicylate and menthol. Methyl salicylate may have an indication for chronic pain in this context. Per MTUS p105, "Recommended. Topical salicylate (e.g., Ben-Gay, methyl salicylate) is significantly better than placebo in chronic pain. (Mason-BMJ, 2004). "However, the CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of menthol. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status equivalent to "not recommended". Since menthol is not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication

change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually.