

Case Number:	CM15-0049762		
Date Assigned:	03/23/2015	Date of Injury:	02/29/2012
Decision Date:	05/06/2015	UR Denial Date:	02/13/2015
Priority:	Standard	Application Received:	03/16/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Indiana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 34 year old female, who sustained an industrial injury on 2/29/2012. She reported an initial injury to her right knee while transferring a patient, with subsequent orthopedic complaints regarding her neck, bilateral shoulders, and low back. A right knee arthroscopy in 2011 and a long history of mental health difficulties were noted. The injured worker was diagnosed as having adjustment disorder with mixed features, anxiety and depression, chronic, probable personality traits/disorder, chronic pain, lumbar facet arthropathy, lumbar degenerative disc disease, and myofascial pain. Treatment to date has included diagnostics, medications, bilateral medial branch blocks (with greater than 80% relief noted), and physical therapy. On 1/02/2015, the injured worker complained of low back pain, rated 6-8/10. Physical exam noted tenderness to the bilateral L4-5 region and increased pain with range of motion testing. The treatment plan included Gabapentin, Valium, and Percocet. A progress report, dated 5/20/2014, noted the use of Gabapentin and Diazepam. Pain levels were not documented.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Gabapentin 300mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-epilepsy drug (AEDs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-epilepsy drugs Page(s): 16-22. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, Anti-epilepsy drugs (AEDs) for pain, Gabapentin (Neurontin®).

Decision rationale: The MTUS considers Gabapentin as a first-line treatment for neuropathic pain and effective for the treatment of spinal cord injury, lumbar spinal stenosis, and post op pain. MTUS also recommends a trial of Gabapentin for complex regional pain syndrome. ODG states "Recommended Trial Period: One recommendation for an adequate trial with Gabapentin is three to eight weeks for titration, then one to two weeks at maximum tolerated dosage. (Dworkin, 2003) The patient should be asked at each visit as to whether there has been a change in pain or function. Current consensus based treatment algorithms for diabetic neuropathy suggests that if inadequate control of pain is found, a switch to another first-line drug is recommended." Additionally, ODG states that Gabapentin "has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain." Based on the clinical documentation provided, there is no evidence of neuropathic type pain or radicular pain on exam or subjectively. As such, without any evidence of neuropathic type pain, the medication is not medically necessary.

Flexeril 10mg #30: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Chronic Pain.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine; anti-spasmodics Page(s): 41-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Cyclobenzaprine (Flexeril®), Other Medical Treatment Guideline or Medical Evidence: UpToDate, Flexeril.

Decision rationale: MTUS Chronic Pain Medical Treatment states for Cyclobenzaprine, "Recommended as an option, using a short course of therapy. The effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. (Browning, 2001) Treatment should be brief." The medical documents indicate that patient is far in excess of the initial treatment window and period. Additionally, MTUS outlines that "Relief of pain with the use of medications is generally temporary, and measures of the lasting benefit from this modality should include evaluating the effect of pain relief in relationship to improvements in function and increased activity. Before prescribing any medication for pain the following should occur: (1) determine the aim of use of the medication; (2) determine the potential benefits and adverse effects; (3) determine the patient's preference. Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be

recorded. (Mens, 2005)" Uptodate "flexeril" also recommends "Do not use longer than 2-3 weeks". Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine.ODG states regarding cyclobenzaprine, "Recommended as an option, using a short course of therapy. The addition of cyclobenzaprine to other agents is not recommended." Several other pain medications are being requested, along with cyclobenzaprine, which ODG recommends against. As such, the request is not medically necessary.