

Case Number:	CM15-0050122		
Date Assigned:	03/23/2015	Date of Injury:	07/02/2014
Decision Date:	05/06/2015	UR Denial Date:	02/26/2015
Priority:	Standard	Application Received:	03/17/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 66-year-old male who sustained an industrial injury on July 2, 2014. He has reported injury to the cervical and lumbar spine and has been diagnosed with cervical spine sprain/strain and lumbar spine sprain/strain. Treatment has included medications and physical therapy. Recent progress noted he suffered an injury to his cervical spine and lumbar spine secondary to a fall. The treatment request included Tramadol, cyclobenzaprine, and topical cream.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Tramadol 50mg quantity 90: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol Page(s): 113.

Decision rationale: According to MTUS guidelines, Ultram (Tramadol) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework. There is no clear documentation of pain and functional improvement with previous use of the Tramadol. There is no clear documentation of continuous compliance of the patient with his medications. There is no documentation of the medical necessity of Tramadol over NSAID. Therefore, the prescription of Tramadol 50 mg #90 is not medically necessary.

Cyclobenzaprine 7.5mg quantity 90: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants Page(s): 63.

Decision rationale: According to MTUS guidelines, Cyclobenzaprine a non sedating muscle relaxants is recommended with caution as a second line option for short term treatment of acute exacerbations in patients with chronic spasm and pain. Efficacy appears to diminish over time and prolonged use may cause dependence. The guidelines do not recommend being used form more than 2-3 weeks. The patient in this case does not have clear recent evidence of spasm and the prolonged use of Cyclobenzaprine is not justified. Therefore, the request for Cyclobenzaprine 7.5mg #90 is not medically necessary.

Topical Cream: Lidocaine 2%, 30gm: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111.

Decision rationale: According to MTUS, in Chronic Pain Medical Treatment, guidelines section Topical Analgesics (page 111); topical analgesics are largely experimental in use with few randomized controlled trials to determine efficacy or safety. Many agents are combined to other pain medications for pain control. There is limited research to support the use of many of these agents. Furthermore, according to MTUS guidelines, any compounded product that contains at least one drug or drug class that is not recommended is not recommended. According to the patient file, there is no documentation of failure of first line therapies or functional improvement with previous use of Lidocain cream. There is no evidence of neuropathic origin of the patient pain. Therefore the prescription of Topical Cream: Lidocaine 2%, 30gm is not medically necessary.