

<b>Case Number:</b>	CM15-0050624		
<b>Date Assigned:</b>	03/24/2015	<b>Date of Injury:</b>	05/23/1997
<b>Decision Date:</b>	05/01/2015	<b>UR Denial Date:</b>	03/13/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	03/17/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: New York, Tennessee  
 Certification(s)/Specialty: Emergency Medicine

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51 year old male, who sustained an industrial injury on 5/23/1997. The mechanism of injury was not provided for review. The injured worker was diagnosed as having lumbosacral neuritis, left lower extremity healing ulcer, depression, insomnia and chronic left leg pain. There is no record of a recent diagnostic study. Treatment to date has included therapy and medication management. Currently, the injured worker complains of left leg pain. In a progress note dated 2/10/2015, the treating physician is requesting Norco and Neurontin.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Norco 10/325 mg, 180 count:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 11, 74-96.

**Decision rationale:** Norco is the compounded medication containing hydrocodone and acetaminophen. Chronic Pain Medical Treatment Guidelines state that opioids are not recommended as a first line therapy. Opioid should be part of a treatment plan specific for the patient and should follow criteria for use. Criteria for use include establishment of a treatment plan, determination if pain is nociceptive or neuropathic, failure of pain relief with non-opioid analgesics, setting of specific functional goals, and opioid contract with agreement for random drug testing. If analgesia is not obtained, opioids should be discontinued. The patient should be screened for likelihood that he or she could be weaned from the opioids if there is no improvement in pain or function. It is recommended for short term use if first-line options, such as acetaminophen or NSAIDs have failed. Opioids may be a safer choice for patients with cardiac and renal disease than antidepressants or anticonvulsants. Acetaminophen is recommended for treatment of chronic pain & acute exacerbations of chronic pain. Acetaminophen overdose is a well-known cause of acute liver failure. Hepatotoxicity from therapeutic doses is unusual. Renal insufficiency occurs in 1 to 2% of patients with overdose. The recommended dose for mild to moderate pain is 650 to 1000 mg orally every 4 hours with a maximum of 4 g/day. In this case the patient has been receiving Norco since at least September 2014 and has not obtained analgesia. In addition there is no documentation that the patient has signed an opioid contract or is participating in urine drug testing. Criteria for long-term opioid use have not been met. The request is not medically necessary.

**Neurontin 300 mg, sixty count:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 18-19.

**Decision rationale:** Neurontin is the anti-epileptic medication, gabapentin. Gabapentin has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain and has FDA approval for treatment of post-herpetic neuralgia. Gabapentin appears to be effective in reducing abnormal hypersensitivity, to have anti-anxiety effects, and may be beneficial as a sleep aid. Gabapentin has a favorable side-effect profile, few clinically significant drug-drug interactions and is generally well tolerated; however, common side effects include dizziness, somnolence, confusion, ataxia, peripheral edema, dry mouth, and weight gain. It has been recommended for the treatment of pain from spinal cord injury, fibromyalgia, lumbar spinal stenosis, and chronic regional pain syndrome. Recommended trial period is three to eight weeks for titration, then one to two weeks at maximum tolerated dosage. If inadequate control of pain is found, a switch to another first-line drug is recommended. In this case, the patient has been treated with neurontin since at least September 2014 and has not obtained analgesia. The medication should be discontinued if there is no functional benefit. The request is not medically necessary.