

Case Number:	CM15-0050752		
Date Assigned:	03/24/2015	Date of Injury:	07/17/2012
Decision Date:	05/08/2015	UR Denial Date:	03/06/2015
Priority:	Standard	Application Received:	03/17/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Indiana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 58 year old male, who sustained an industrial injury on 7/17/2012. He reported back pain. Diagnoses have included left shoulder impingement syndrome with full thickness tearing of the left rotator cuff tendon with retraction with infraspinatus tendinitis and AC joint arthritis. Treatment to date has included diagnostic studies including magnetic resonance imaging (MRI), physical therapy, analgesics and anti-inflammatory medications. Per the Secondary Treating Physician's Initial Orthopaedic Surgical Consultation Report dated 6/26/2014, the injured worker reported constant, severe left shoulder pain. Physical examination revealed marked tenderness over the left shoulder AC joint region and over the anterior aspect of the left shoulder. Range of motion was reduced. There was a strongly positive Impingement sign, and Neer and Hawkin's tests. Yergason's test, Drop arm sign, Codman's test and abrasion test for left shoulder rotator cuff tendinopathy were positive. The plan of care included surgical intervention and medications and authorization was requested for Tramadol 15mg #60, Naproxen 550mg #90 and Pantoprazole 20mg #90.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Tramadol 150mg #60: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol; Opioids Page(s): 74-123. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic) - Medications for acute pain (analgesics), Tramadol (Ultram®).

Decision rationale: Ultram is the brand name version of tramadol, which is classified as central acting synthetic opioids. MTUS states regarding tramadol that a therapeutic trial of opioids should not be employed until the patient has failed a trial of non-opioid analgesics. Before initiating therapy, the patient should set goals, and the continued use of opioids should be contingent on meeting these goals. ODG further states, Tramadol is not recommended as a first-line oral analgesic because of its inferior efficacy to a combination of Hydrocodone/acetaminophen. The treating physician did not provide sufficient documentation that the patient has failed a trial of non-opioid analgesics at the time of prescription or in subsequent medical notes. Additionally, no documentation was provided which discussed the setting of goals for the use of tramadol prior to the initiation of this medication. As such, the request is not medically necessary.

Naproxen 550mg #90: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Naproxen, NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: MTUS specifies four recommendations regarding NSAID use: 1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. 2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP. 3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. 4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. Additionally, the treating physician does not document failure of primary (Tylenol) treatment. Progress notes do not indicate how long the patient has been on naproxen, but the MTUS guidelines recommend against long-term use. As such, the request is not medically necessary.

Pantoprazole 20mg #90: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS; GI protection Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDS, GI symptoms & cardiovascular risk.

Decision rationale: MTUS states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." ODG states, if a PPI is used, omeprazole OTC tablets or lansoprazole 24 HR OTC are recommended for an equivalent clinical efficacy and significant cost savings. Products in this drug class have demonstrated equivalent clinical efficacy and safety at comparable doses, including esomeprazole (Nexium), lansoprazole (Prevacid), omeprazole (Prilosec), pantoprazole (Protonix), dexlansoprazole (Dexilant), and rabeprazole (Aciphex). (Shi, 2008) A trial of omeprazole or lansoprazole is recommended before Nexium therapy. The other PPIs, Protonix, Dexilant, and Aciphex, should also be second-line. According to the latest AHRQ Comparative Effectiveness Research, all of the commercially available PPIs appeared to be similarly effective. (AHRQ, 2011) The patient does not meet the age recommendations for increased GI risk. The medical documents provided establish the patient has experienced GI discomfort, but is nonspecific and does not indicate history of peptic ulcer, GI bleeding or perforation. Medical records do not indicate that the patient is on ASA, corticosteroids, and/or an anticoagulant; or high dose/multiple NSAID. Additionally per guidelines, Pantoprazole is considered second line therapy and the treating physician has not provided detailed documentation of a failed trial of omeprazole and/or lansoprazole. As such, the request for Pantoprazole is not medically necessary.