

Case Number:	CM15-0050989		
Date Assigned:	03/24/2015	Date of Injury:	12/06/1998
Decision Date:	05/01/2015	UR Denial Date:	03/12/2015
Priority:	Standard	Application Received:	03/17/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 56-year-old female who sustained an industrial injury on 12/6/1998. Her diagnoses, and/or impressions, include right shoulder pain; left wrist pain; left thumb pain; and trigger finger. No recent magnetic resonance imaging studies were noted. Her treatments have included left thumb and left little finger A1 pulley release (10/31/14), desensitization exercises and medication management. In the progress note dated 12/22/2014, she reported for re-evaluation following the A1 pulley release, the finger remained sore and no further triggering was noted. The physician's requests for treatment included Naprosyn for pain and Prilosec for non-steroidal anti-inflammatory induced dyspepsia, from Naprosyn.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Naprosyn 550mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs) Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Naproxen, NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: MTUS specifies four recommendations regarding NSAID use: 1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. 2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP. 3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. 4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. Additionally, the treating physician does not document failure of primary (Tylenol) treatment. Progress notes do not indicate how long the patient has been on naproxen, but the MTUS guidelines recommend against long-term use. The treating physician has not provided documentation of objective functional improvement to warrant the continued use of this medication. As such, the request for Naprosyn 550mg #60 is not medically necessary.

Prilosec 20mg #30, 1 every day, refills: 3: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risks with precautions.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: MTUS states "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." The medical documents provided do not establish the patient has having documented GI bleeding/perforation/peptic ulcer or other GI risk factors as outlined in MTUS. Additionally, there is no evidence provided to indicate the patient suffers from

dyspepsia because of the present medication regimen. As such, the request for Prilosec 20mg #30, 1 every day, refills: 3 is not medically necessary.