

Case Number:	CM15-0051779		
Date Assigned:	03/25/2015	Date of Injury:	01/11/2001
Decision Date:	05/01/2015	UR Denial Date:	03/12/2015
Priority:	Standard	Application Received:	03/19/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The 61 year old male injured worker suffered an industrial injury on 01/11/2001. The diagnoses included lumbosacral neuritis, lumbago, and lumbar spinal stenosis without claudication, joint pain in forearm, depression and joint pain in pelvis. The injured worker had been treated with medications. On 3/4/2015, the treating provider reported left foot burning pain and left lower leg pain 7 to 8/10 increased significantly with the decrease of OxyContin. He walks with a limp utilizing a walker with wheels and armrest. The treatment plan included Xanax and Norco.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Xanax XR 2mg (Alprazolam) #150: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

Decision rationale: According to MTUS guidelines, benzodiazepines are not recommended for long-term use for pain management because of unproven long-term efficacy and because of the risk of dependence. Most guidelines limit their use to 4 weeks. Xanax was used for a long time without any evidence of efficacy. Therefore, the use of Xanax XR 2mg #150 is not medically necessary.

Norco 10/325mg #180: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria for use of opioids Page(s): 76-79.

Decision rationale: According to MTUS guidelines, Norco (Hydrocodone/Acetaminophen) is a synthetic opioid indicated for the pain management but not recommended as a first line oral analgesic. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules:”(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework.” According to the patient file, there is no objective documentation of pain and functional improvement to justify continuous use of Norco. Norco was used for longtime without documentation of functional improvement or evidence of return to work or improvement of activity of daily living. Therefore, the prescription of Norco 10/325mg #180 is not medically necessary.