

Case Number:	CM15-0053705		
Date Assigned:	03/27/2015	Date of Injury:	01/07/2015
Decision Date:	05/01/2015	UR Denial Date:	02/23/2015
Priority:	Standard	Application Received:	03/20/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: North Carolina
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 33 year old male, who sustained an industrial injury on 1/7/15. He reported pain in the lower back related to a motor vehicle accident. The injured worker was diagnosed as having lumbar sprain and intercostal strain. Treatment to date has included acupuncture, chiropractic treatments, lumbar CT and pain medications. As of the PR2 dated 2/19/15, the injured worker reports 9/10 pain in the lower back that radiates to the bilateral lower extremities. The treating physician noted tenderness in the lumbar paravertebral muscles and restricted range of motion. The treating physician requested an open MRI of the lumbar spine and a home IF unit.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

MRI Lumbar Spine (open): Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303-304.

Decision rationale: The ACOEM chapter on low back complaints and special diagnostic studies states: Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false-positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computed tomography [CT] for bony structures). Relying solely on imaging studies to evaluate the source of low back and related symptoms carries a significant risk of diagnostic confusion (false positive test results) because of the possibility of identifying a finding that was present before symptoms began and therefore has no temporal association with the symptoms. Techniques vary in their abilities to define abnormalities (Table 12-7). Imaging studies should be reserved for cases in which surgery is considered or red-flag diagnoses are being evaluated. Because the overall false-positive rate is 30% for imaging studies in patients over age 30 who do not have symptoms, the risk of diagnostic confusion is great. There is no recorded presence of emerging red flags on the physical exam. There is evidence of nerve compromise on physical exam but there is not mention of consideration for surgery or complete failure of conservative therapy. For these reasons, criteria for imaging as defined above per the ACOEM have not been met. Therefore, the request is not medically necessary.

Home Interferential Unit: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines TENS (transcutaneous electrical nerve stimulation).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines interferential therapy Page(s): 118-119.

Decision rationale: The California medical treatment guidelines section on ICS therapy states: Not recommended as an isolated intervention. There is no quality evidence of effectiveness except in conjunction with recommended treatments, including return to work, exercise and medications, and limited evidence of improvement on those recommended treatments alone. The randomized trials that have evaluated the effectiveness of this treatment have included studies for back pain, jaw pain, soft tissue shoulder pain, cervical neck pain and post-operative knee pain. (Van der Heijden, 1999)(Werner, 1999) (Hurley, 2001) (Hou, 2002) (Jarit, 2003) (Hurley, 2004) (CTAF, 2005) (Burch, 2008) The findings from these trials were either negative or non-interpretable for recommendation due to poor study design and/or methodologic issues. In addition, although proposed for treatment in general for soft tissue injury or for enhancing wound or fracture healing, there is insufficient literature to support Interferential current stimulation for treatment of these conditions. There are no standardized protocols for the use of interferential therapy; and the therapy may vary according to the frequency of stimulation, the pulse duration, treatment time, and electrode-placement technique.

Two recent randomized double-blind controlled trials suggested that ICS and horizontal therapy (HT) were effective in alleviating pain and disability in patients with chronic low back pain compared to placebo at 14 weeks, but not at 2 weeks. The placebo effect was remarkable at the beginning of the treatment but it tended to vanish within a couple of weeks. The studies suggested that their main limitation was the heterogeneity of the low back pain subjects, with the interventions performing much better for back pain due to previous multiple vertebral osteoporotic fractures, and further studies are necessary to determine effectiveness in low back pain from other causes. (Zambito, 2006) (Zambito, 2007) A recent industry-sponsored study in the Knee Chapter concluded that interferential current therapy plus patterned muscle stimulation (using the RS-4i Stimulator) has the potential to be a more effective treatment modality than conventional low-current TENS for osteoarthritis of the knee. (Burch, 2008) This recent RCT found that either electroacupuncture or interferential electrotherapy, in combination with shoulder exercises, is equally effective in treating frozen shoulder patients. It should be noted that this study only showed the combined treatment effects with exercise as compared to no treatment, so the entire positive effect could have been due to the use of exercise alone. (Cheing, 2008) See also Sympathetic therapy. See also TENS, chronic pain. While not recommended as an isolated intervention, Patient selection criteria if Interferential stimulation is to be used anyway: Possibly appropriate for the following conditions if it has documented and proven to be effective as directed or applied by the physician or a provider licensed to provide physical medicine: Pain is ineffectively controlled due to diminished effectiveness of medications; or Pain is ineffectively controlled with medications due to side effects; or History of substance abuse; or Significant pain from postoperative conditions limits the ability to perform exercise programs/physical therapy treatment; or Unresponsive to conservative measures (e.g., repositioning, heat/ice, etc.). If those criteria are met, then a one-month trial may be appropriate to permit the physician and physical medicine provider to study the effects and benefits. There should be evidence of increased functional improvement, less reported pain and evidence of medication reduction. 'jacket' should not be certified until after the one-month trial and only with documentation that the individual cannot apply the stimulation pads alone or with the help of another available person. The criteria as set forth above per the California MTUS have not been met. In addition, ICS is only initially approved for a one-month trial period. Therefore, the request is not medically necessary.