

<b>Case Number:</b>	CM15-0053794		
<b>Date Assigned:</b>	03/27/2015	<b>Date of Injury:</b>	07/23/2011
<b>Decision Date:</b>	05/04/2015	<b>UR Denial Date:</b>	03/06/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	03/20/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Maryland, Texas, Virginia

Certification(s)/Specialty: Internal Medicine, Allergy and Immunology, Rheumatology

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 62 year old female who sustained an industrial injury on July 23, 2011. She has reported low back pain and has been diagnosed with low back pain, facet pain, sacroiliitis, and possibility of lumbar radiculopathy. Treatment has included modified duty, medications, and a TENS unit. Currently the injured worker reports a constant achy pain with intermittent sharp shooting and stabbing pain radiating to the left thigh. The treatment request included gabapentin and tramadol.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Gabapentin 100mg 1-2 tablets oral at bedtime #60, refill 3:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Anti epilepsy drugs (AEDs) Page(s): 16-19.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Anti-Epilepsy Drugs Page(s): 16-22. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chronic Pain, Anti-epilepsy drugs (AEDs) for pain, Gabapentin (Neurontin).

**Decision rationale:** The MTUS considers Gabapentin as a first-line treatment for neuropathic pain and effective for the treatment of spinal cord injury, lumbar spinal stenosis, and post op pain. MTUS also recommends a trial of Gabapentin for complex regional pain syndrome. ODG states "Recommended Trial Period: One recommendation for an adequate trial with Gabapentin is three to eight weeks for titration, then one to two weeks at maximum tolerated dosage. (Dworkin, 2003) The patient should be asked at each visit as to whether there has been a change in pain or function. Current consensus based treatment algorithms for diabetic neuropathy suggests that if inadequate control of pain is found, a switch to another first-line drug is recommended". Additionally, ODG states that Gabapentin "has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain". Based on the clinical documentation provided, there is no evidence of neuropathic type pain or radicular pain on exam or subjectively. As such, without any evidence of neuropathic type pain, the medication is not medically necessary.

**Tramadol 50mg 1 tablet oral every 12 hours #60 refill 3:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioid, on going management Page(s): 78-80.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Tramadol, Ultram Page(s): 74-96, 113, 123. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic) - Medications for acute pain (analgesics), Tramadol (Ultram).

**Decision rationale:** Tramadol is classified as a central acting synthetic opioids. MTUS states regarding tramadol that "A therapeutic trial of opioids should not be employed until the patient has failed a trial of non-opioid analgesics. Before initiating therapy, the patient should set goals, and the continued use of opioids should be contingent on meeting these goals". ODG further states, "Tramadol is not recommended as a first-line oral analgesic because of its inferior efficacy to a combination of Hydrocodone/ acetaminophen". The treating physician did not provide sufficient documentation that the patient has failed a trial of non-opioid analgesics at the time of prescription or in subsequent medical notes. Additionally, no documentation was provided which discussed the setting of goals for the use of tramadol prior to the initiation of this medication. MTUS states that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life". The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for Tramadol 50mg 1 tablet oral every 12 hours #60 refills 3 is not medically necessary.

