

Case Number:	CM15-0054723		
Date Assigned:	03/30/2015	Date of Injury:	10/27/2008
Decision Date:	05/01/2015	UR Denial Date:	03/12/2015
Priority:	Standard	Application Received:	03/23/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: New Jersey

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 49 year old male who sustained an industrial injury on 10/27/08. The injured worker reported symptoms in the back and lower extremities. The injured worker was diagnosed as having nerve compression and L4-L5 disc compression. Treatments to date have included nonsteroidal anti-inflammatory drugs, topical creams, status post fusion, physical therapy, epidural steroid injection, status post laminotomy, and aqua therapy. Currently, the injured worker complains of pain in the lower back with radiation to the lower extremities. The plan of care was for medication prescriptions and a follow up appointment at a later date.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Percocet 10/325mg #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 78.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 78-96.

Decision rationale: The MTUS Chronic Pain Medical Treatment Guidelines state that opioids may be considered for moderate to severe chronic pain as a secondary treatment, but require that for continued opioid use, there is to be ongoing review and documentation of pain relief, functional status, appropriate medication use with implementation of a signed opioid contract, drug screening (when appropriate), review of non-opioid means of pain control, using the lowest possible dose, making sure prescriptions are from a single practitioner and pharmacy, and side effects, as well as consultation with pain specialist if after 3 months unsuccessful with opioid use, all in order to improve function as criteria necessary to support the medical necessity of opioids. Long-term use and continuation of opioids requires this comprehensive review with documentation to justify continuation. In the case of this worker, there was insufficient documentation to show this full review regarding Percocet use was completed. There was no report of functional gains and measurable pain level reduction directly related to Percocet. Therefore, the request for Percocet will be considered medically unnecessary at this time.

Omeprazole: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk Page(s): 68-69.

Decision rationale: The MTUS Guidelines state that to warrant using a proton pump inhibitor (PPI) in conjunction with an NSAID, the patient would need to display intermediate or high risk for developing a gastrointestinal event such as those older than 65 years old, those with a history of peptic ulcer, GI bleeding, or perforation, or those taking concurrently aspirin, corticosteroids, and/or an anticoagulant, or those taking a high dose or multiple NSAIDs. In the case of this worker, there was no documentation suggesting this worker was at an elevated risk for gastrointestinal events to warrant ongoing omeprazole use. Also, the request was missing the dose and number of pills. Therefore, the Omeprazole will be considered medically unnecessary.