

Case Number:	CM15-0055232		
Date Assigned:	04/17/2015	Date of Injury:	12/16/2012
Decision Date:	05/18/2015	UR Denial Date:	03/03/2015
Priority:	Standard	Application Received:	03/19/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, District of Columbia, Maryland
 Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 30 year old female, who sustained an industrial injury on December 16, 2012. She reported back pain radiating to the legs. The injured worker was diagnosed as having lumbosacral radiculitis and degenerative disc disease. Treatment to date has included diagnostic studies, physical therapy, steroid injections, medications and work restrictions. Currently, the injured worker complains of back pain radiating to the legs. The injured worker reported an industrial injury in 2012, resulting in the above noted pain. She was treated conservatively without complete resolution of the pain. Evaluation on December 16, 2012, revealed continued pain as noted. It was noted she required medications to maintain function. A pain management consultation and Protonix were requested.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Protonix 20mg 1-2 every morning, #60: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter, NSAIDs, GI symptoms and cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI Symptoms & Cardiovascular Risk Page(s): 68. Decision based on Non-MTUS Citation ODG, Pain, Proton Pump Inhibitors.

Decision rationale: In the treatment of dyspepsia secondary to NSAID therapy, the MTUS recommends stopping the NSAID, switching to a different NSAID, or considering the use of an H2-receptor antagonist or a PPI. The MTUS Chronic Pain Medical Treatment Guidelines recommend the use of proton pump inhibitors in conjunction with NSAIDs in situations in which the patient is at risk for gastrointestinal events including: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). CPMTG guidelines further specify. Recommendations: -Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g., ibuprofen, naproxen, etc.). Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Patients at high risk of gastrointestinal events with cardiovascular disease: If GI risk is high the suggestion is for a low-dose Cox-2 plus low dose Aspirin (for cardioprotection) and a PPI. If cardiovascular risk is greater than GI risk the suggestion is naproxyn plus low-dose aspirin plus a PPI. (Laine, 2006) (Scholmerich, 2006) (Nielsen, 2006) (Chan, 2004) (Gold, 2007) (Laine, 2007)" Per ODG TWC, "many prescribers believe that this class of drugs is innocuous, but much information is available to demonstrate otherwise. A trial of omeprazole or lansoprazole is recommended before Nexium therapy. The other PPIs, Protonix, Dexilant, and Aciphex, should also be second-line." As there is no documentation of peptic ulcer, GI bleeding or perforation, or cardiovascular disease in the records available for my review, the injured worker's risk for gastrointestinal events is low, as such, medical necessity cannot be affirmed. Furthermore, as noted per the guidelines, Protonix is a second-line medication. The medical records do not establish whether the patient has failed attempts at first line PPIs, such as omeprazole or lansoprazole, which should be considered prior to prescribing a second line PPI such as Protonix. The request is not medically necessary.

Pain management consult: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation ACOEM, page 127 Official Disability Guidelines (ODG), Low Back Chapter.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 2 General Approach to Initial Assessment and Documentation Page(s): 27.

Decision rationale: The California MTUS Guidelines recommend a consultation to aid with diagnosis/prognosis and therapeutic management, recommend referrals to other specialist if a diagnosis is uncertain or exceedingly complex when there are psychosocial factors present, or when, a plan or course of care may benefit from additional expertise. The medical necessity of

the requested referral has not been sufficiently established by the documentation available for my review. The documentation does not specify what the pain management consult will address. The request is not medically necessary.