

<b>Case Number:</b>	CM15-0055241		
<b>Date Assigned:</b>	03/30/2015	<b>Date of Injury:</b>	12/15/2001
<b>Decision Date:</b>	05/11/2015	<b>UR Denial Date:</b>	03/19/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	03/23/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 62 year old male sustained an industrial injury to the neck, back, right upper extremity and shoulder on 12/15/01. Previous treatment included magnetic resonance imaging, electromyography, lumbar surgery, physical therapy, chiropractic therapy, acupuncture, psychological care, epidural steroid injections, sacroiliac joint blocks, trigger point injections, transcutaneous electrical nerve stimulator unit and meds. In a PR-2 dated 3/5/15, the injured worker complained of pain 9/10 on the visual analog scale to the cervical spine, right shoulder and right low back and buttock area with radiation to the right lower extremity. The injured worker reported that he had been walking more lately and now had right shin pain. The injured worker reported stopping Norco due to nausea. Current diagnoses included lumbar spine radiculopathy, failed lumbar back surgery syndrome, cervical spine radiculopathy, right shoulder impingement syndrome, myofascial pain syndrome, chronic pain and right ankle/foot joint pain. The treatment plan included discontinuing Norco, a prescription for Tramadol and physical therapy.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Tramadol HCL 50mg, #30 with 1 refill: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Tramadol (Ultram).

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-97.

**Decision rationale:** According to the MTUS Chronic Pain Medical Treatment Guidelines section on Opioids, On-Going Management, p 74-97, (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the injured worker's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the injured worker's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain injured workers on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or no adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the injured worker should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or uninjured worker treatment with issues of abuse, addiction, or poor pain control.(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).(g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Additionally, the MTUS states that continued use of opioids requires (a) the injured worker has returned to work, (b) the injured worker has improved functioning and pain. There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

**Norco 5/325mg #30:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, specific drug list, Criteria for use of Opioids.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids  
Page(s): 74-97.

**Decision rationale:** According to the MTUS Chronic Pain Medical Treatment Guidelines section on Opioids, On-Going Management, p 74-97, (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the injured worker's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the injured worker's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain injured workers on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or no adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the injured worker should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or uninjured worker treatment with issues of abuse, addiction, or poor pain control.(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).(g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Additionally, the MTUS states that continued use of opioids requires (a) the injured worker has returned to work, (b) the injured worker has improved functioning and pain. There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

**1 urine drug screen:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain (Chronic).

**MAXIMUS guideline:** The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Drug Testing.

**Decision rationale:** ODG, Pain (Chronic), Drug Testing Recommended as a tool to monitor compliance with prescribed substances, identify use of undisclosed substances, and uncover diversion of prescribed substances. The test should be used in conjunction with other clinical information when decisions are to be made to continue, adjust or discontinue treatment. This information includes clinical observation, results of addiction screening, pill counts, and prescription drug monitoring reports. The prescribing clinician should also pay close attention to information provided by family members, other providers and pharmacy personnel. The frequency of urine drug testing may be dictated by state and local laws. Indications for UDT: At the onset of treatment: (1) UDT is recommended at the onset of treatment of a new injured worker who is already receiving a controlled substance or when chronic opioid management is considered. Urine drug testing is not generally recommended in acute treatment settings (i.e. when opioids are required for nociceptive pain). (2) In cases in which the injured worker asks for a specific drug. This is particularly the case if this drug has high abuse potential; the injured worker refuses other drug treatment and/or changes in scheduled drugs, or refuses generic drug substitution. (3) If the injured worker has a positive or "at risk" addiction screen on evaluation. This may also include evidence of a history of comorbid psychiatric disorder such as depression, anxiety, bipolar disorder, and/or personality disorder. See Opioids, screening tests for risk of addiction & misuse. (4) If aberrant behavior or misuse is suspected and/or detected. See Opioids, indicators for addiction & misuse. Ongoing monitoring: (1) If an injured worker has evidence of a "high risk" of addiction (including evidence of a comorbid psychiatric disorder (such as depression, anxiety, attention-deficit disorder, obsessive-compulsive disorder, bipolar disorder, and/or schizophrenia), has a history of aberrant behavior, personal or family history of substance dependence (addiction), or a personal history of sexual or physical trauma, ongoing urine drug testing is indicated as an adjunct to monitoring along with clinical exams and pill counts. See Opioids, tools for risk stratification & monitoring. (2) If dose increases are not decreasing pain and increasing function, consideration of UDT should be made to aid in evaluating medication compliance and adherence. According to the documents available for review, the injured worker meets none of the aforementioned MTUS criteria for the use of urine drug testing. Therefore at this time the requirements for treatment have not been met, and medical necessity has not been established.

**1 gym membership:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Low back - Lumbar & Thoracic (Acute & Chronic).

**MAXIMUS guideline:** Decision based on MTUS ACOEM Chapter 3 Initial Approaches to Treatment.

**Decision rationale:** The ACOEM Chapter 3 on Treatment indicates that specialized treatments or referrals require a rationale for their use. According to the documents available for review, there is no rationale provided to support the request of a gym membership. Therefore at this time the requirements for treatment have not been met, and medical necessity has not been established.

## **16 physical therapy sessions: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Physical medicine guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98.

**Decision rationale:** Physical Medicine is recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the injured worker) can provide short term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Injured workers are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Injured worker-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of injured workers with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks According to the documents available for review, the current request of the total number of sessions is in contrast to the guidelines as set forth in the MTUS. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.