

Case Number:	CM15-0055835		
Date Assigned:	04/01/2015	Date of Injury:	07/23/2003
Decision Date:	05/06/2015	UR Denial Date:	03/13/2015
Priority:	Standard	Application Received:	03/24/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: District of Columbia, Virginia
 Certification(s)/Specialty: Internal Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 54 year old, female who sustained a work related injury on 7/23/03. The diagnoses have included bilateral lateral epicondylitis, status post bilateral elbow surgery and chronic pain syndrome. Treatments have included acupuncture, medications, an MRI and an electrodiagnostic study. In an Initial Evaluation Report dated 1/21/15, the injured worker complains of pain in left shoulder, both elbows, both wrists, lower back, both knees and left foot. She describes the pain as frequent and severe. She rates the pain a 9-10/10. She states her activities are restricted due to pain. The treatment plan is to request authorization for a multidisciplinary evaluation.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Multidisciplinary evaluation: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines functional restoration programs (FPRs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines 9792
 Page(s): 31-33.

Decision rationale: Per MTUS: Predictors of success and failure: As noted, one of the criticisms of interdisciplinary/multidisciplinary rehabilitation programs is the lack of an appropriate screening tool to help to determine who will most benefit from this treatment. Retrospective research has examined decreased rates of completion of functional restoration programs, and there is ongoing research to evaluate screening tools prior to entry. (Gatchel, 2006) The following variables have been found to be negative predictors of efficacy of treatment with the programs as well as negative predictors of completion of the programs: (1) a negative relationship with the employer/supervisor; (2) poor work adjustment and satisfaction; (3) a negative outlook about future employment; (4) high levels of psychosocial distress (higher pretreatment levels of depression, pain and disability); (5) involvement in financial disability disputes; (6) greater rates of smoking; (7) duration of pre-referral disability time; (8) prevalence of opioid use; and (9) pretreatment levels of pain. (Linton, 2001) (Bendix, 1998) (McGeary, 2006) (McGeary, 2004) (Gatchel, 2005) Multidisciplinary treatment strategies are effective for patients with chronic low back pain (CLBP) in all stages of chronicity and should not only be given to those with lower grades of CLBP, according to the results of a prospective longitudinal clinical study reported in the December 15 issue of Spine. (Buchner, 2007) See also Chronic pain programs, early intervention; Chronic pain programs, intensity; Chronic pain programs, opioids; and Functional restoration programs. Criteria for the general use of multidisciplinary pain management programs: Outpatient pain rehabilitation programs may be considered medically necessary when all of the following criteria are met: (1) An adequate and thorough evaluation has been made, including baseline functional testing so follow-up with the same test can note functional improvement; (2) Previous methods of treating chronic pain have been unsuccessful and there is an absence of other options likely to result in significant clinical improvement; (3) The patient has a significant loss of ability to function independently resulting from the chronic pain; (4) The patient is not a candidate where surgery or other treatments would clearly be warranted (if a goal of treatment is to prevent or avoid controversial or optional surgery, a trial of 10 visits may be implemented to assess whether surgery may be avoided); (5) The patient exhibits motivation to change, and is willing to forgo secondary gains, including disability payments to effect this change; & (6) Negative predictors of success above have been addressed. Integrative summary reports that include treatment goals, progress assessment and stage of treatment, must be made available upon request and at least on a bi-weekly basis during the course of the treatment program. Treatment is not suggested for longer than 2 weeks without evidence of demonstrated efficacy as documented by subjective and objective gains. (Note: Patients may get worse before they get better. For example, objective gains may be moving joints that are stiff from lack of use, resulting in increased subjective pain.) However, it is also not suggested that a continuous course of treatment be interrupted at two weeks solely to document these gains, if there are preliminary indications that these gains are being made on a concurrent basis. Total treatment duration should generally not exceed 20 full-day sessions (or the equivalent in part-day sessions if required by part-time work, transportation, childcare, or comorbidities). (Sanders, 2005) Treatment duration in excess of 20 sessions requires a clear rationale for the specified extension and reasonable goals to be achieved. Longer durations require individualized care plans and proven outcomes, and should be based on chronicity of disability and other known risk factors for loss of function. Inpatient pain rehabilitation programs: These programs typically consist of more intensive functional rehabilitation and medical care than their outpatient counterparts. They may be appropriate for patients who: (1) don't have the minimal functional capacity to participate effectively in an outpatient program; (2) have medical conditions that require more intensive oversight; (3) are receiving large amounts of medications necessitating medication weaning or detoxification; or (4) have complex medical or psychological diagnosis that benefit from more intensive observation and/or additional consultation during the rehabilitation process. (Keel, 1998) (Kool, 2005) (Buchner, 2006) (Kool, 2007) As with outpatient pain rehabilitation programs, the most effective programs combine intensive, daily bio-psychosocial rehabilitation

with a functional restoration approach. (BlueCross BlueShield, 2004) (Aetna, 2006) See Functional restoration programs. Based on the clinical documentation provided, it appears that multiple therapies had been tried on this patient. Despite these, the patient had ongoing pain issues. The patient was referred to an orthopedist but had not yet had a formal evaluation. Per cited guidelines, a multidisciplinary approach would be appropriate.