

Case Number:	CM15-0056711		
Date Assigned:	04/01/2015	Date of Injury:	06/17/2013
Decision Date:	05/04/2015	UR Denial Date:	02/27/2015
Priority:	Standard	Application Received:	03/25/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 48-year-old male with an industrial injury dated June 17, 2013. The injured worker diagnoses include lateral collateral ligament laxity status post reefing, low back pain, status post total knee arthroplasty, tibial plateau fracture of the left knee, insomnia, reactive depression, and left lower leg vascular injury. He has been treated with diagnostic studies, home exercises, physical therapy, prescribed medications, knee surgery and periodic follow up visits. According to the progress note dated February 24, 2015, the injured worker reported left knee pain. The treating physician noted left knee tenderness at medial compartment greater than the lateral compartment and mid lateral instability. The treating physician prescribed physical therapy for the left knee and Tramadol 50mg now under review.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Physical therapy 2 times a week for 4 weeks for the left knee: Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines: Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active, self directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks. Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks. Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The requested amount of physical therapy is within guideline recommendations and therefore is medically necessary.

Trazadone 50mg #60 with 1 refill: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation ODG, insomnia.

Decision rationale: The California MTUS and the ACOEM do not specifically address this medication. Per the official disability guidelines recommend pharmacological agents for insomnia only is used after careful evaluation of potential causes of sleep disturbance. Primary insomnia is usually addressed pharmacologically. Secondary insomnia may be treated with pharmacological and/or psychological measures. Pharmacological treatment consists of four main categories: Benzodiazepines, Non-benzodiazepines, Melatonin and melatonin receptor

agonists and over the counter medications. Sedating antidepressants have also been used to treat insomnia however, there is less evidence to support their use for insomnia, but they may be option in patients with coexisting depression. Trazadone is a sedating antidepressant. The patient does not have a coexisting diagnosis of depression. Therefore, criteria for its use have not been met and the request is not medically necessary.