

Case Number:	CM15-0057073		
Date Assigned:	04/01/2015	Date of Injury:	04/26/2012
Decision Date:	05/07/2015	UR Denial Date:	02/26/2015
Priority:	Standard	Application Received:	03/25/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: California, Arizona, Maryland
Certification(s)/Specialty: Psychiatry

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 34 year old female, who sustained an industrial injury on 4/26/2012. She reported the base of a bed fell onto the right foot. This acute event was treated with Vicodin, a brace, crutches and activity modification. The medical records indicated the onset of psychological symptoms followed secondary to chronic pain and stress. Diagnoses include major depressive disorder, and pain disorder associated with both medical and psychological factors. Treatments to date include medication therapy, activity modification, and psychotherapy. Currently, she reported some improvement in anxiety and irritability. On 2/20/15, the physical examination documented the affect was good and appropriate. The diagnoses was depression, major/single. The plan of care included continuation of medication therapy including Trazodone and Lexapro as ordered.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Trazodone 10mg 1 tab q HS PRN #30 with 2 refills: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 15 Stress Related Conditions Page(s): 394-404. Decision based on Non-MTUS Citation Official Disability

Guidelines (ODG), Treatment Index, 11th Edition (web), 2014, Mental Illness and Stress, Antidepressants for treatment of MDD (major depressive disorder).

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Mental Illness & Stress Trazodone (Desyrel).

Decision rationale: ODG states that Trazodone: Recommended as an option for insomnia, only for patients with potentially coexisting mild psychiatric symptoms such as depression or anxiety. See also Insomnia treatment, where it says there is limited evidence to support its use for insomnia, but it may be an option in patients with coexisting depression. See also Fibromyalgia in the Pain Chapter, where trazodone was used successfully in fibromyalgia. Trazodone was approved in 1982 for the treatment of depression. It is unrelated to tricyclic or tetracyclic antidepressants and has some action as an anxiolytic. Off-label uses include alcoholism, anxiety, insomnia, and panic disorder. Although approved to treat depression, the American Psychiatric Association notes that it is not typically used for major depressive disorder. Over the period 1987 through 1996, prescribing trazodone for depression decreased throughout the decade, while off-label use of the drug for insomnia increased steadily until it was the most frequently prescribed insomnia agent. To date, there has been only one randomized, double blind, placebo-controlled trial studying trazodone in primary insomnia. It was observed that relative to placebo, patients reported significant improvement in subjective sleep latency, sleep duration, wake time after sleep onset, and sleep quality with trazodone and zolpidem during week one, but during week two the trazodone group did not differ significantly from the placebo group whereas the zolpidem group demonstrated significant improvement compared to placebo for sleep latency and sleep duration. (Walsh, 1998) The AHRQ Comparative Effectiveness Research on insomnia concludes that trazodone is equal to zolpidem. (AHRQ, 2008) Evidence for the off-label use of trazodone for treatment of insomnia is weak. The current recommendation is to utilize a combined pharmacologic and psychological and behavior treatment when primary insomnia is diagnosed. Also worth noting, there has been no dose-finding study performed to assess the dose of trazodone for insomnia in non-depressed patients. Other pharmacologic therapies should be recommended for primary insomnia before considering trazodone, especially if the insomnia is not accompanied by comorbid depression or recurrent treatment failure. There is no clear-cut evidence to recommend trazodone first line to treat primary insomnia. (Mendelson, 2005) The treating provider's progress notes indicate that Trazodone is being prescribed in this case for treatment of insomnia coexisting with depression. Thus, the request for Trazodone 10mg 1 tab q HS PRN #30 with 2 refills is medically necessary.

Lexapro 10mg 1 tab daily #30 with 2 refills: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 15 Stress Related Conditions. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Treatment Index, 11th Edition (web), 2014, Mental Illness and Stress, Antidepressants for treatment of MDD (major depressive disorder).

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Chapter: Stress & Mental Illness Topic: Antidepressants for treatment of MDD (major depressive disorder).

Decision rationale: ODG states "MDD (major depressive disorder) treatment, severe presentations; The American Psychiatric Association strongly recommends anti-depressant medications for severe presentations of MDD, unless electroconvulsive therapy (ECT) is being planned. (American Psychiatric Association, 2006) Many treatment plans start with a category of medication called selective serotonin reuptake inhibitors (SSRIs), because of demonstrated effectiveness and less severe side effects." The request for Lexapro 10mg 1 tab daily #30 with 2 refills is medically necessary for the ongoing symptoms of Major Depressive Disorder in this case.