

Case Number:	CM15-0057460		
Date Assigned:	04/02/2015	Date of Injury:	03/13/2001
Decision Date:	05/05/2015	UR Denial Date:	02/26/2015
Priority:	Standard	Application Received:	03/26/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 48 year old female, who sustained an industrial injury on March 13, 2001. She reported neck pain, left shoulder pain, low back pain, bilateral lower extremity pain and left knee pain. The injured worker was diagnosed as having herniated lumbar disc, lumbar radiculopathy, impingement syndrome of the bilateral shoulders, cervical disc displacement, status post cervical fusion, cervical radiculopathy and left knee internal derangement. Treatment to date has included radiographic imaging, diagnostic studies, surgical intervention of the cervical spine, conservative therapies, medications and work restrictions. Currently, the injured worker complains of neck pain, left shoulder pain, low back pain, bilateral lower extremity pain and left knee pain. The injured worker reported an industrial injury in 2001, resulting in the above noted pain. She was treated conservatively and surgically without complete resolution of the pain. Evaluation on January 23, 2015, revealed continued pain as noted. Medications were adjusted and renewed.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

6 NORCO: STRENGTH: 10/325MG; QUANTITY: 120; REFILLS: UNSPECIFIED; TAKEN BY MOUTH, 1 TABLET EVERY 4 HOURS AS NEEDED: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Shoulder, Pain, Opioids.

Decision rationale: ODG does not recommend the use of opioids for neck, low back, and shoulder pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. Additionally, medical documents indicate that the patient has been on Norco in excess of the recommended 2-week limit. As such, the request for 6 NORCO: STRENGTH: 10/325MG; QUANTITY: 120; REFILLS: UNSPECIFIED; TAKEN BY MOUTH, 1 TABLET EVERY 4 HOURS AS NEEDED is not medically necessary.

FLEXERIL: STRENGTH 10 MG; QUANTITY: 90; REFILLS: UNSPECIFIED; TAKEN BY MOUTH, 1 TABLET THREE TIMES A DAY: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxers.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cyclobenzaprine, Medications for chronic pain, Antispasmodics Page(s): 41-42, 60-61, 64-66. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Cyclobenzaprine (Flexeril[®] 1/2) and Other Medical Treatment Guidelines UpToDate, Flexeril.

Decision rationale: MTUS Chronic Pain Medical Treatment states for Cyclobenzaprine, "Recommended as an option, using a short course of therapy. The effect is greatest in the first 4 days of treatment, suggesting that shorter courses may be better. (Browning, 2001) Treatment should be brief." The medical documents indicate that patient is far in excess of the initial treatment window and period. Additionally, MTUS outlines that "Relief of pain with the use of medications is generally temporary, and measures of the lasting benefit from this modality should include evaluating the effect of pain relief in relationship to improvements in function and increased activity. Before prescribing any medication for pain the following should occur: (1) determine the aim of use of the medication; (2) determine the potential benefits and adverse effects; (3) determine the patient's preference. Only one medication should be given at a time,

and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) Uptodate "flexeril" also recommends "Do not use longer than 2-3 weeks". Medical documents do not fully detail the components outlined in the guidelines above and do not establish the need for long term/chronic usage of cyclobenzaprine. ODG states regarding cyclobenzaprine, "Recommended as an option, using a short course of therapy. The addition of cyclobenzaprine to other agents is not recommended." Several other pain medications are being requested, along with cyclobenzaprine, which ODG recommends against. As such, the request for FLEXERIL: STRENGTH 10 MG; QUANTITY: 90; REFILLS: UNSPECIFIED; TAKEN BY MOUTH, 1 TABLET THREE TIMES A DAY is not medically necessary.

ULTRAM: STRENGTH: 50 MG; QUANTITY: 90; REFILLS: UNSPECIFIED; TAKEN BY MOUTH, 1 CAPSULE, THREE TIMES A DAY: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Tramadol, Ultram Page(s): 74-96, 113, 123. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic) - Medications for acute pain (analgesics), Tramadol (Ultram ½).

Decision rationale: Tramadol is classified as a central acting synthetic opioids. MTUS states regarding tramadol that "A therapeutic trial of opioids should not be employed until the patient has failed a trial of non-opioid analgesics. Before initiating therapy, the patient should set goals, and the continued use of opioids should be contingent on meeting these goals." ODG further states, "Tramadol is not recommended as a first-line oral analgesic because of its inferior efficacy to a combination of Hydrocodone/ acetaminophen." The treating physician did not provide sufficient documentation that the patient has failed a trial of non-opioid analgesics at the time of prescription or in subsequent medical notes. Additionally, no documentation was provided which discussed the setting of goals for the use of tramadol prior to the initiation of this medication. MTUS states that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the request for ULTRAM: STRENGTH: 50 MG; QUANTITY: 90; REFILLS: UNSPECIFIED; TAKEN BY MOUTH, 1 CAPSULE, THREE TIMES A DAY is not medically necessary.