

<b>Case Number:</b>	CM15-0057463		
<b>Date Assigned:</b>	04/02/2015	<b>Date of Injury:</b>	03/30/2011
<b>Decision Date:</b>	06/29/2015	<b>UR Denial Date:</b>	03/17/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	03/26/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Maryland, Texas, Virginia

Certification(s)/Specialty: Internal Medicine, Allergy and Immunology, Rheumatology

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 46 year old male who sustained an industrial injury on 3/30/11. The injured worker was diagnosed as having lower back pain and deep vein thrombosis on chronic Coumadin therapy. Currently, the injured worker was with complaints of lower extremity weakness. Previous treatments included status post laminotomy and physical therapy. Previous diagnostic studies included a computed tomography angiogram, magnetic resonance imaging and ultrasound venous Doppler study. The plan of care was for an epidural steroid injection.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Epidural Steroid injection at Left L3-4 (specified hospital): Upheld**

**Claims Administrator guideline:** The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation CA MTUS: page 46, 2010 Revision, Web Edition.

**MAXIMUS guideline:** Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 287-315, Chronic Pain Treatment Guidelines Epidural steroid injections (ESIs) Page(s): 46. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Low Back - Lumbar & Thoracic (Acute & Chronic), Epidural steroid injections (ESIs), therapeutic.

**Decision rationale:** MTUS Chronic pain medical treatment guidelines state that epidural steroid injections are "Recommended as an option for treatment of radicular pain (defined as pain in dermatomal distribution with corroborative findings of radiculopathy). Epidural steroid injection can offer short term pain relief and use should be in conjunction with other rehab efforts, including continuing a home exercise program." There were no medical documents provided to conclude that other rehab efforts or home exercise program is ongoing. Additionally, no objective findings were documented to specify the dermatomal distribution of pain. MTUS further defines the criteria for epidural steroid injections to include: 1) Radiculopathy must be documented by physical examination and corroborated by imaging studies and/or electrodiagnostic testing. 2) Initially unresponsive to conservative treatment (exercises, physical methods, NSAIDs and muscle relaxants). 3) Injections should be performed using fluoroscopy (live x-ray) for guidance. 4) If used for diagnostic purposes, a maximum of two injections should be performed. A second block is not recommended if there is inadequate response to the first block. Diagnostic blocks should be at an interval of at least one to two weeks between injections. 5) No more than two nerve root levels should be injected using transforaminal blocks. 6) No more than one interlaminar level should be injected at one session. 7) In the therapeutic phase, repeat blocks should be based on continued objective documented pain and functional improvement, including at least 50% pain relief with associated reduction of medication use for six to eight weeks, with a general recommendation of no more than 4 blocks per region per year. (Manchikanti, 2003) (CMS, 2004) (Boswell, 2007)8) Current research does not support a 'series-of-three' injections in either the diagnostic or therapeutic phase. We recommend no more than 2 ESI injections. It is unclear if radiculopathy is demonstrated with imaging studies as these are not provided. The physical exam does not demonstrate radiculopathy. The patient is taking multiple medications, but the progress reports do not document how long the patient has been on these medications and the 'unresponsiveness' to the medications. Additionally, treatment notes do not indicate if other conservative treatments were tried and failed (exercises, physical therapy, etc). As such, the request for Epidural steroid injection at left L3-4 (specified hospital) is not medically necessary.