

Case Number:	CM15-0058151		
Date Assigned:	04/02/2015	Date of Injury:	08/25/2011
Decision Date:	06/12/2015	UR Denial Date:	03/03/2015
Priority:	Standard	Application Received:	03/26/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Arizona, Michigan

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 27 year old male, who sustained an industrial injury on 8/25/2011. He reported pain after bending over to pick up a heavy panel of glass. The injured worker was diagnosed as having lumbar disc herniation, bilateral lower extremity radicular pain and neuroforaminal stenosis bilaterally. There is no record of a recent diagnostic study. Treatment to date has included medication management. In a progress note dated 2/23/2015, the injured worker complains of lumbar spine pain, which improves with rest and medication. The treating physician is requesting urine drug screen, Motrin, Prilosec and Flurbiprofen/Lidocaine cream.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Urine Toxicology Screen: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Drug testing Page(s): 43. Decision based on Non-MTUS Citation Official Disability Guidelines TWC, Urine Drug Testing (UDT).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Drug Testing Page(s): 43. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic)/Urine Drug testing.

Decision rationale: Per the MTUS, Drug testing is recommended as an option, using a urine drug screen to assess for the use or the presence of illegal drugs before a therapeutic trial of opioids, during ongoing management and to avoid misuse/addiction. Per the ODG, frequency of urine drug testing should be based on documented evidence of risk stratification including use of a testing instrument. A review of the injured workers medical records did not reveal documentation that the injured worker is on opioid therapy, an intention to begin opioid therapy or risk stratification and without this information the request is not medically necessary.

Flurbiprofen/Lidocaine Cream 20%/5% 180gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

Decision rationale: Per the MTUS, topical analgesics are recommended as an option, they are largely experimental in use with few randomized controlled trials to determine efficacy or safety. They are primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. Many agents are compounded as monotherapy or in combination for pain control, any compounded product that contains at least one drug or drug class that is not recommended is not recommended. A review of the injured workers medical records that are available to me does not show a trial of recommended first line agents that have failed therefore the request for Flurbiprofen/Lidocaine Cream 20%/5% 180gm is not medically necessary.

Motrin 800mg #60: Overturned

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 67-68.

Decision rationale: Per the MTUS, NSAIDs are recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that

long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. A review of the injured workers medical records that are available to me reveal documentation of the injured workers improvement in pain and function with the use of Motrin therefore the continued use of Motrin 800 mg # 60 is medically necessary.

Prilosec 20mg #60: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms and cardiovascular risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic)/Proton Pump Inhibitors (PPIs).

Decision rationale: Per the MTUS, Clinicians should weigh the indications for NSAIDs against both GI and cardiovascular risk factors according to specific criteria listed in the MTUS and a selection should be made based on these criteria 1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Per the ODG, PPIs are Recommended for patients at risk for gastrointestinal events. Prilosec (omeprazole), Prevacid (lansoprazole) and Nexium (esomeprazole magnesium) are PPIs. Healing doses of PPIs are more effective than all other therapies, although there is an increase in overall adverse effects compared to placebo. Nexium and Prilosec are very similar molecules. (Donnellan, 2010) In this RCT omeprazole provided a statistically significantly greater acid control than lansoprazole. (Miner, 2010) In general, the use of a PPI should be limited to the recognized indications and used at the lowest dose for the shortest possible amount of time. PPIs are highly effective for their approved indications, including preventing gastric ulcers induced by NSAIDs. Studies suggest, however, that nearly half of all PPI prescriptions are used for unapproved indications or no indications at all. Many prescribers believe that this class of drugs is innocuous, but much information is available to demonstrate otherwise. Products in this drug class have demonstrated equivalent clinical efficacy and safety at comparable doses, including esomeprazole (Nexium), lansoprazole (Prevacid), omeprazole (Prilosec), pantoprazole (Protonix), dexlansoprazole (Dexilant), and rabeprazole (Aciphex). (Shi, 2008) A trial of omeprazole or lansoprazole had been recommended before prescription Nexium therapy (before it went OTC). The other PPIs, Protonix, Dexilant, and Aciphex, should be second-line. According to the latest AHRQ Comparative Effectiveness Research, all of the commercially available PPIs appeared to be similarly effective. (AHRQ, 2011). A review of the injured workers medical records that are available to me do not reveal any gastrointestinal complaints or history and the injured worker does not meet the guideline criteria for increased risk, therefore the request for Prilosec 20mg #60 is not medically necessary.