

Case Number:	CM15-0060205		
Date Assigned:	04/06/2015	Date of Injury:	05/13/2011
Decision Date:	05/05/2015	UR Denial Date:	03/12/2015
Priority:	Standard	Application Received:	03/30/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is 41 year old male, who sustained an industrial injury on 05/13/2011. He reported low back pain and bilateral wrist pain. He was diagnosed with wrist and lumbosacral strains. Treatment to date has included x-rays, MRI, physical therapy, chiropractic care, acupuncture, medications and TENS unit. The injured worker complained of constant lower back pain with radiation to the left leg, aching in both shoulder girdles and at the back of his neck and constant throbbing, burning pain of his arms, wrists and hands. Currently under review is the request for Ultram, Terocin patches and electrodes #4 for the transcutaneous nerve stimulation unit.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Ultram 50mg, #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines When to continue opioids. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG): Pain Chapter, Opioids, criteria for use; Opioids for chronic pain.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids
Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids; (a) If the patient has returned to work. (b) If the patient has improved functioning and pain (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented improvement in VAS scores. There are also no objective measurements of improvement in function. Therefore criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

Terocin patch 4%, #30 patches: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines topical analgesics Page(s): 111-113.

Decision rationale: The California chronic pain medical treatment guidelines section on topical analgesics states: Recommended as an option as indicated below. Largely experimental in use with few randomized controlled trials to determine efficacy or safety. Primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. (Namaka, 2004) These agents are applied locally to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. (Colombo, 2006) Many agents are compounded as monotherapy or in combination for pain control (including NSAIDs, opioids, capsaicin, local anesthetics, antidepressants, glutamate receptor antagonists, adrenergic receptor agonist, adenosine, cannabinoids, cholinergic receptor agonists, agonists, prostanoids, bradykinin, adenosine triphosphate, biogenic amines, and nerve growth factor). (Argoff, 2006) There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The requested medication contains multiple ingredients, which are not indicated per the California MTUS for topical analgesic use. Therefore the request is not medically necessary.

Durable medical equipment (DME) electrodes #4, for the transcutaneous electrical nerve stimulation (TENS) unit: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines TENS, chronic pain (transcutaneous electrical nerve stimulation).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines TENS Page(s): 114.

Decision rationale: The California chronic pain medical treatment guidelines section on transcutaneous electrical nerve stimulation states: TENS, chronic pain (transcutaneous electrical nerve stimulation) not recommended as a primary treatment modality, but a one-month home-based TENS trial may be considered as a noninvasive conservative option, if used as an adjunct to a program of evidence-based functional restoration, for the conditions described below. While TENS may reflect the long-standing accepted standard of care within many medical communities, the results of studies are inconclusive; the published trials do not provide information on the stimulation parameters which are most likely to provide optimum pain relief, nor do they answer questions about long-term effectiveness. (Carroll-Cochrane, 2001) Several published evidence-based assessments of transcutaneous electrical nerve stimulation (TENS) have found that evidence is lacking concerning effectiveness. One problem with current studies is that many only evaluated single-dose treatment, which may not reflect the use of this modality in a clinical setting. Other problems include statistical methodology, small sample size, influence of placebo effect, and difficulty comparing the different outcomes that were measured. This treatment option is recommended as an adjunct to a program of evidence based functional restoration. The patient has documented long-term use of TENS unit with benefit. Therefore, the request is medically necessary.