MAXIMUS Federal Services

Case Number:	CM15-0080508		
Date Assigned:	05/01/2015	Date of Injury:	01/20/2000
Decision Date:	06/01/2015	UR Denial Date:	04/14/2015
Priority:	Standard	Application Received:	04/27/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
State(s) of Licensure: North Carolina
Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 66-year-old male, who sustained an industrial injury on January 20, 2000. He has reported neck pain, back pain, and leg pain. Diagnoses have included chronic pain, lumbar spine radiculitis, and reflex sympathetic dystrophy. Treatment to date has included medications and a spinal cord stimulator. A progress note dated April 7, 2015 indicates a chief complaint of back pain, buttock pain, and leg and foot pain. The injured worker also complained of increasing leg weakness. The treating physician documented a plan of care that included medications and sympathetic blocks.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg quantity 180: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000)(d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain dairy that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore criteria for the ongoing use of opioids have not been met and the request is not medically necessary.

Bilateral L2 paravertebral sympathetic blocks: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Regional Sympathetic Blocks.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines CRPS Page(s): 39.

Decision rationale: The California MTUS section on CRPS and sympathetic blocks states: CRPS, sympathetic and epidural blocks recommended only as indicated below, for a limited role, primarily for diagnosis of sympathetically mediated pain and as an adjunct to facilitate physical therapy. Detailed information about stellate ganglion blocks, thoracic sympathetic blocks, and lumbar sympathetic blocks is found in Regional sympathetic blocks. Recommendations for the use of sympathetic blocks are listed below. They are recommended for a limited role, primarily for diagnosis of sympathetically mediated pain and as an adjunct to facilitate physical therapy. It should be noted that sympathetic blocks are not specific for CRPS. See Sympathetically maintained pain (SMP). Repeated blocks are only recommended if continued improvement is observed. Systematic reviews reveal a paucity of published evidence supporting the use of local anesthetic sympathetic blocks for the treatment of CRPS and usefulness remains controversial. Less than 1/3 of patients with CRPS are likely to respond to sympathetic blockade. No controlled trials have shown any significant benefit from sympathetic blockade. (Varrassi, 2006) (Cepeda, 2005) (Hartrick, 2004) (Grabow, 2005) (Cepeda, 2002) (Forouzanfar, 2002) (Sharma, 2006) Predictors of poor response: Long duration of symptoms prior to intervention; Elevated anxiety levels; Poor coping skills; Litigation. (Hartrick, 2004) (Nelson, 2006) Alternatives to regional sympathetic blocks: may be necessary when there is evidence of coagulopathy, systemic infection, and/or post-surgical changes. These include peripheral nerve and plexus blocks and epidural administration of local anesthetics. Mixed conduction blocks (central neural blocks): suggested when analgesia is insufficient by pharmacologic means to support physical therapy: (1) Implanted catheters at the brachial or lumbosacral plexus: allows for 1 to 2 weeks of therapy. Side effects include technical failure and infection; & (2) Epidural tunneled catheters: allows for long-term therapy: Side effects: same as above. The provided documentation for review does not meet criteria as stated above for sympathetic blocks in the treatment of CRPS and therefore the request is not medically necessary.

Right C6-7 sympathetic stellate ganglion block: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Regional Sympathetic Blocks.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines CRPS Page(s): 39.

Decision rationale: The California MTUS section on CRPS and sympathetic blocks states: CRPS, sympathetic and epidural blocks recommended only as indicated below, for a limited role, primarily for diagnosis of sympathetically mediated pain and as an adjunct to facilitate physical therapy. Detailed information about stellate ganglion blocks, thoracic sympathetic blocks, and lumbar sympathetic blocks is found in Regional sympathetic blocks. Recommendations for the use of sympathetic blocks are listed below. They are recommended for a limited role, primarily for diagnosis of sympathetically mediated pain and as an adjunct to facilitate physical therapy. It should be noted that sympathetic blocks are not specific for CRPS. See Sympathetically maintained pain (SMP). Repeated blocks are only recommended if continued improvement is observed. Systematic reviews reveal a paucity of published evidence supporting the use of local anesthetic sympathetic blocks for the treatment of CRPS and usefulness remains controversial. Less than 1/3 of patients with CRPS are likely to respond to sympathetic blockade. No controlled trials have shown any significant benefit from sympathetic blockade. (Varrassi, 2006) (Cepeda, 2005) (Hartrick, 2004) (Grabow, 2005) (Cepeda, 2002) (Forouzanfar, 2002) (Sharma, 2006) Predictors of poor response: Long duration of symptoms prior to intervention; Elevated anxiety levels; Poor coping skills; Litigation. (Hartrick, 2004)

(Nelson, 2006) Alternatives to regional sympathetic blocks: may be necessary when there is evidence of coagulopathy, systemic infection, and/or post-surgical changes. These include peripheral nerve and plexus blocks and epidural administration of local anesthetics. Mixed conduction blocks (central neural blocks): suggested when analgesia is insufficient by pharmacologic means to support physical therapy: (1) Implanted catheters at the brachial or lumbosacral plexus: allows for 1 to 2 weeks of therapy. Side effects include technical failure and infection; & (2) Epidural tunneled catheters: allows for long-term therapy: Side effects: same as above. The provided documentation for review does not meet criteria as stated above for sympathetic blocks in the treatment of CRPS and therefore the request is not medically necessary.