

Case Number:	CM15-0080623		
Date Assigned:	05/01/2015	Date of Injury:	03/09/2013
Decision Date:	06/02/2015	UR Denial Date:	04/16/2015
Priority:	Standard	Application Received:	04/27/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 42 year old female, who sustained an industrial injury on 03/09/2013. She has reported injury to the neck, right shoulder, and back. The diagnoses have included cervicgia/neck pain; shoulder, joint pain; and backache, unspecified. Treatment to date has included medications, diagnostic studies, TENS (transcutaneous electrical nerve stimulation) unit, home exercises, and surgical intervention. Medications have included Gabapentin, Omeprazole, Lidopro Cream, and Docuprene. A progress note from the treating physician, dated 04/03/2015, documented a follow-up visit with the injured worker. Currently, the injured worker complains of chronic neck pain; bilateral trapezii muscle spasm; right shoulder and mid back pain; upper extremity pain has increased and changed to constant as well as increased symptoms; and sleep issues which are affecting her mood. Objective findings included tenderness to palpation in the right shoulder, trapezius muscle, and cervical paraspinal muscles; and decreased range of motion to the cervical spine and the right shoulder. The treatment plan has included the request for Omeprazole 20mg #60; Gabapentin 500mg #160; and Lidopro Cream 121gm.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Omeprazole 20mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Nsaids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAID Page(s): 68-70.

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy and proton pump inhibitors (PPI) states: Recommend with precautions as indicated below. Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastro duodenal lesions. Recommendations-Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g,ibuprofen, naproxen, etc.). Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastro-intestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. There is no documentation provided that places this patient at intermediate or high risk that would justify the use of a PPI. There is no mention of current gastrointestinal or cardiovascular disease. For these reasons the criteria set forth above per the California MTUS for the use of this medication has not been met. Therefore, the request is not medically necessary.

Gabapentin 500mg #160: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-Epilepsy.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines gabapentin Page(s): 18.

Decision rationale: The California chronic pain medical treatment guidelines section on Neurontin states: Gabapentin (Neurontin, Gabarone, generic available) has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain. (Backonja, 2002) (ICSI, 2007) (Knotkova, 2007) (Eisenberg, 2007) (Attal, 2006) This RCT concluded that gabapentin monotherapy appears to be efficacious for the treatment of pain and sleep interference associated with diabetic peripheral neuropathy and exhibits positive effects on mood and quality of life. (Backonja, 1998) It has been given FDA approval for treatment of post-herpetic neuralgia. The number needed to treat (NNT) for overall neuropathic pain is 4. It has a more favorable side-effect profile than Carbamazepine, with a number needed to harm of 2.5. (Wiffen2-Cochrane, 2005) (Zaremba,2006) Gabapentin in combination with morphine has been studied for treatment of diabetic neuropathy and postherpetic neuralgia. When used in combination the maximum tolerated dosage of both drugs was lower than when each was used as a single agent and better

analgesia occurred at lower doses of each. (Gilron-NEJM, 2005) Recommendations involving combination therapy require further study. The requested medication is a first line agent to treatment neuropathic pain. The patient does not have a diagnosis of neuropathic pain. Therefore, the request is not medically necessary.

Lidopro Cream 121gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines topical analgesics Page(s): 111-113.

Decision rationale: The California chronic pain medical treatment guidelines section on topical analgesics states: Recommended as an option as indicated below. Largely experimental in use with few randomized controlled trials to determine efficacy or safety. Primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. (Namaka, 2004) These agents are applied locally to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. (Colombo, 2006) Many agents are compounded as monotherapy or in combination for pain control (including NSAIDs, opioids, capsaicin, local anesthetics, antidepressants, glutamate receptor antagonists, & #945;-adrenergic receptor agonist, adenosine, cannabinoids, cholinergic receptor agonists, & #947; agonists, prostanoids, bradykinin, adenosine triphosphate, biogenic amines, and nerve growth factor). (Argoff, 2006) There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The requested medication contains ingredients, which are not indicated per the California MTUS for topical analgesic use. Therefore, the request is not medically necessary.