

<b>Case Number:</b>	CM15-0082281		
<b>Date Assigned:</b>	05/04/2015	<b>Date of Injury:</b>	04/27/2009
<b>Decision Date:</b>	06/04/2015	<b>UR Denial Date:</b>	04/20/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	04/29/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: California  
 Certification(s)/Specialty: Physical Medicine & Rehabilitation

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 60 year old male, who sustained an industrial injury on 4/27/09. The injured worker has complaints of left shoulder tenderness. The diagnoses have included status-post right shoulder surgery and pain in joint, shoulder region. Treatment to date has included right shoulder surgery in January 2015; physical therapy; transcutaneous electrical nerve stimulation unit; home exercise program; hydrocodone; xanax and omeprazole. The request was for hydrocodone 10mg, #90 and xanax 0.5mg, #60.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Hydrocodone 10mg, #90:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not cite any medical evidence for its decision.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96.

**Decision rationale:** Per the MTUS Guidelines cited, opioid use in the setting of chronic, non-malignant, or neuropathic pain is controversial. Patients on opioids should be routinely monitored for signs of impairment and use of opioids in patients with chronic pain should be reserved for those with improved functional outcomes attributable to their use, in the context of an overall approach to pain management that also includes non-opioid analgesics, adjuvant therapies, psychological support, and active treatments (e.g., exercise). Submitted documents show no evidence that the treating physician is prescribing opioids in accordance to change in pain relief, functional goals with demonstrated improvement in daily activities, decreased in medical utilization or change in functional status. There is no evidence presented of random drug testing or utilization of pain contract to adequately monitor for narcotic safety, efficacy, and compliance. The MTUS provides requirements of the treating physician to assess and document for functional improvement with treatment intervention and maintenance of function that would otherwise deteriorate if not supported. From the submitted reports, there is no demonstrated evidence of specific functional benefit derived from the continuing use of opioids with persistent severe pain for this chronic injury without acute flare, new injury, or progressive deterioration. The Hydrocodone 10mg, #90 is not medically necessary and appropriate.

**Xanax 0.5mg, #60:** Upheld

**Claims Administrator guideline:** The Claims Administrator did not cite any medical evidence for its decision.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

**Decision rationale:** Xanax Tablets (alprazolam) is indicated for the management of anxiety disorder. Anxiety or tension associated with the stress of everyday life usually does not require treatment with an anxiolytic. Alprazolam is an anti-anxiety medication in the benzodiazepine family which inhibits many of the activities of the brain as it is believed that excessive activity in the brain may lead to anxiety or other psychiatric disorders. Per the Chronic Pain Treatment Guidelines, benzodiazepines are not recommended for long-term use because long-term efficacy is unproven and there is a risk of dependence. Most guidelines limit use to 4 weeks as chronic benzodiazepines are the treatment of choice in very few conditions and tolerance to hypnotic effects develops rapidly. The Xanax 0.5mg, #60 is not medically necessary and appropriate.