

Case Number:	CM15-0083467		
Date Assigned:	05/04/2015	Date of Injury:	09/27/2002
Decision Date:	06/05/2015	UR Denial Date:	03/30/2015
Priority:	Standard	Application Received:	04/28/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker (IW) is a 60 year old male who sustained an industrial injury on 09/27/2002. He reported pain in his right shoulder. The injured worker was diagnosed as having impingement syndrome of the shoulder on the right status post two surgical interventions, and chronic pain syndrome with associated weight gain of 30 pounds, and issues with sleep, stress, and depression, approved as being work-related. Treatment to date has included oral pain medications, psychiatric evaluation, treatment with antidepressants, and cognitive behavioral therapy. Currently, the injured worker complains of his right shoulder and his psyche issues. He has limitation with forceful activities, pushing, pulling and overhead work. Sleep is affected, and he has spasms. His physician describes that he has issues with sleep, stress, and depression, and that there is tenderness along the shoulder girdle musculature and the rotator cuff. Abduction is 150 degrees. Impingement sign is negative. He has weakness to resisted function. A request for authorization is placed for one prescription of Naproxen 500mg #60, and one prescription of Pantoprazole 20mg #60. On 03/30/2015 the utilization reviews organization non-certified both requests citing MTUS Chronic Pain.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

1 prescription of Naproxen 500mg #60: Overturned

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAID Page(s): 68-70.

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy states: Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain: Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. This medication is recommended for the shortest period of time and at the lowest dose possible. The dosing of this medication is within the California MTUS guideline recommendations. The definition of shortest period possible is not clearly defined in the California MTUS. Therefore the request is medically necessary.

1 prescription of Pantoprazole 20mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAID Page(s): 68-70.

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy and proton pump inhibitors (PPI) states: Recommend with precautions as indicated below. Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2)

history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDs to develop gastro duodenal lesions. Recommendations:-Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 µg four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. There is no documentation provided that places this patient at intermediate or high risk that would justify the use of a PPI. There is no mention of current gastrointestinal or cardiovascular disease. For these reasons the criteria set forth above per the California MTUS for the use of this medication has not been met. Therefore the request is not medically necessary.