

Case Number:	CM15-0084310		
Date Assigned:	05/06/2015	Date of Injury:	11/08/2013
Decision Date:	06/11/2015	UR Denial Date:	04/14/2015
Priority:	Standard	Application Received:	05/01/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: New York, Tennessee

Certification(s)/Specialty: Emergency Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 37-year-old male sustained an industrial injury on 1/10/13. He subsequently reported back pain. Diagnoses include post-laminectomy syndrome. Treatments to date include x-ray and MRI testing, surgery, physical therapy and prescription pain medications. The injured worker continues to experience low back pain with radiation to the left lower extremity. Upon examination, tenderness is present in the paraspinal muscles of the lower lumbar spine, range of motion is significantly decreased, straight leg raise test was positive on the left, negative on the right. A request for Percocet medication was made by the treating physician.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Percocet 10/325mg, #90: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, specific drug list - Oxycodone/acetaminophen; Criteria for use of Opioids; Weaning of Medications.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 11, 74-96.

Decision rationale: Percocet 10/325 is compounded medication containing oxycodone/acetaminophen. Chronic Pain Medical Treatment Guidelines state that opioids are not recommended as a first line therapy. Opioid should be part of a treatment plan specific for the patient and should follow criteria for use. Criteria for use include establishment of a treatment plan, determination if pain is nociceptive or neuropathic, failure of pain relief with non-opioid analgesics, setting of specific functional goals, and opioid contract with agreement for random drug testing. If analgesia is not obtained, opioids should be discontinued. The patient should be screened for likelihood that he or she could be weaned from the opioids if there is no improvement in pain or function. It is recommended for short-term use if first-line options, such as acetaminophen or NSAIDs have failed. Opioids are considered a second-line treatment for several reasons: (1) head-to-head comparisons have found that opioids produce more side effects than TCAs and gabapentin; (2) long-term safety has not been systematically studied; (3) long term use may result in immunological and endocrine problems (including hypogonadism); (4) treatment may be associated with hyperalgesia; & (5) opioid use is associated with misuse/abuse. Opioids may be a safer choice for patients with cardiac and renal disease than antidepressants or anticonvulsants. Acetaminophen is recommended for treatment of chronic pain & acute exacerbations of chronic pain. Acetaminophen overdose is a well-known cause of acute liver failure. Hepatotoxicity from therapeutic doses is unusual. Renal insufficiency occurs in 1 to 2% of patients with overdose. The recommended dose for mild to moderate pain is 650 to 1000 mg orally every 4 hours with a maximum of 4 g/day. In this case the patient has been receiving opioid medication since at least June 2014 and has not obtained analgesia. In addition, there is no documentation that the patient has signed an opioid contract or is participating in urine drug testing. Criteria for long-term opioid use have not been met. The request is not medically necessary.