

Case Number:	CM15-0087947		
Date Assigned:	05/12/2015	Date of Injury:	09/09/2014
Decision Date:	06/11/2015	UR Denial Date:	04/24/2015
Priority:	Standard	Application Received:	05/07/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 37-year-old male, who sustained an industrial injury on 09/09/2014. Medical records provided by the treating physician did not indicate the injured worker's mechanism of injury. The injured worker was diagnosed as having status post-right knee anterior cruciate ligament (ACL) surgery. Treatment and diagnostic studies to date has included magnetic resonance arthrogram of the right knee, status post diagnostic arthroscopy with subtotal medial meniscectomy, debridement of the anterior cruciate ligament (ACL), and partial synovectomy on 01/06/2015, medication regimen, and as of 04/08/2015 a compilation of sixteen sessions of physical therapy. The progress notes from treating physician are difficult to decipher including the recent progress note from 04/06/2015. In a physical therapy progress note dated 04/03/2015 the therapist reports complaints of intermittent, aching, localized right knee with a pain rating of a 1 out of 10, but the documentation did not indicate if the physical therapy provided any functional improvement. The treating physician requested additional post-operative physical therapy two times four weeks to the right knee for quadriceps strengthening program to increase the range of motion, along with strapping and taping, and a home exercise program.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Additional Post operative Physical Therapy 2 times a week for 4 weeks to right knee:
 Upheld

Claims Administrator guideline: Decision based on MTUS Postsurgical Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine Page(s): 98.

Decision rationale: According to MTUS guidelines, Physical Medicine is recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) There is no documentation of objective findings that support musculoskeletal dysfunction requiring more physical therapy sessions. There is no documentation of the outcome of previous physical therapy sessions and home exercise. There is no documentation supporting additional physical therapy sessions. Therefore Additional Post operative Physical Therapy 2 times a week for 4 weeks to right knee is not medically necessary