

Case Number:	CM15-0089219		
Date Assigned:	05/13/2015	Date of Injury:	08/17/1983
Decision Date:	06/19/2015	UR Denial Date:	05/04/2015
Priority:	Standard	Application Received:	05/08/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: California, Texas, Florida

Certification(s)/Specialty: Anesthesiology, Pain Management, Hospice & Palliative Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 65-year-old male, who sustained an industrial injury on 8/17/1983. The documentation submitted for this review did not include the details regarding the initial injury or the prior treatments to date. Diagnoses include lumbar radiculitis and cervical radiculitis, chronic lumbalgia with lower extremity radicular symptoms and chronic cervicgia. Currently, he complained of pain in the neck, mid back, left arm and low back. All pain was rated 6-7.5/10 VAS. On 4/7/15, the physical examination documented cervical tenderness with positive bilateral compression and Spurling's tests. There was decreased range of motion with pain. The upper extremities were weak with pain during overhead reach. The plan of care included Norco 10/325mg one tablet every six hours #120 and Ultram 50 mg one tablet every four to six hours for pain #90.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines 9792.20 - 9792.26 Page(s): 44, 47, 75-79, 120 of 127.

Decision rationale: Regarding the request for Norco (hydrocodone/acetaminophen), California Pain Medical Treatment Guidelines state that Norco is an opiate pain medication. Due to high abuse potential, close follow-up is recommended with documentation of analgesic effect, objective functional improvement, side effects, and discussion regarding any aberrant use. Guidelines go on to recommend discontinuing opioids if there is no documentation of improved function and pain. Within the documentation available for review, there is no indication that the medication is improving the patient's function or pain (in terms of specific examples of functional improvement and percent reduction in pain or reduced NRS). As such, there is no clear indication for ongoing use of the medication. Opioids should not be abruptly discontinued, but unfortunately, there is no provision to modify the current request to allow tapering. In light of the above issues, the currently requested Norco (hydrocodone/acetaminophen) is not medically necessary.

Ultram 50mg #90 with 1 refill: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines 9792.20 - 9792.26 Page(s): 44, 47, 75-79, 120 of 127.

Decision rationale: Regarding the request for Ultram (tramadol), California Pain Medical Treatment Guidelines state that Ultram is an opiate pain medication. Due to high abuse potential, close follow-up is recommended with documentation of analgesic effect, objective functional improvement, side effects, and discussion regarding any aberrant use. Guidelines go on to recommend discontinuing opioids if there is no documentation of improved function and pain. Within the documentation available for review, there is no indication that the medication is improving the patient's function or pain (in terms of specific examples of functional improvement and percent reduction in pain or reduced NRS). As such, there is no clear indication for ongoing use of the medication. Opioids should not be abruptly discontinued, but fortunately, the last reviewer modified the current request to allow tapering. In light of the above issues, the currently requested Ultram (tramadol) is not medically necessary.