

<b>Case Number:</b>	CM15-0089369		
<b>Date Assigned:</b>	05/13/2015	<b>Date of Injury:</b>	08/13/2004
<b>Decision Date:</b>	06/17/2015	<b>UR Denial Date:</b>	04/10/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	05/08/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 56 year old female, who sustained an industrial injury on 8/13/2004. She reported acute onset low back pain with lifting type activity. Diagnoses include sciatica, chronic myofascial pain syndrome, and mood adjustment disorder secondary to chronic pain. She is status post lumbar fusion in 2007. Treatments to date include activity modification, medication therapy, low back brace, physical therapy, aquatic therapy, and epidural injections. Currently, she complained of back pain with pain down bilateral lower extremities. On 3/12/15, the physical examination documented trigger points palpated in lower shoulder, gluteus medius, and lumbar regions bilaterally. There was decreased range of motion and decreased sensation. Muscle weakness was noted in the hip, knees and ankles. There were positive sacroiliac joint compression tests and positive slump tests. The plan of care included increasing Norco 10/325mg #60 a month to Norco 10/325mg one tablet every six hours #90 a month.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Norco 10/325mg #60:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Criteria for Use.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

**Decision rationale:** ODG does not recommend the use of opioids for neck and low back pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the question for Norco 10/325mg #60 is not medically necessary.

**Norco 10/325mg #90:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Criteria for Use.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Opioids, Pain.

**Decision rationale:** ODG does not recommend the use of opioids for neck and low back pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. As such, the question for Norco 10/325mg #90 is not medically necessary.

