

Case Number:	CM15-0089725		
Date Assigned:	05/14/2015	Date of Injury:	05/28/2014
Decision Date:	09/22/2015	UR Denial Date:	04/16/2015
Priority:	Standard	Application Received:	05/11/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Massachusetts

Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 29-year-old female who sustained an industrial injury on 5/28/2014 resulting in radiating low back pain. She was diagnosed with lumbar sprain and lumbosacral neuritis. Treatment has included physical therapy, epidural steroid injection, and oral and topical medications providing reported minimal, temporary pain relief. The injured worker continues to report low back pain, and bilateral leg pain with numbness and weakness, impacting mobility and sleep. The treating physician's plan of care has included a follow up visit on 5/6/15; and, Anaprox, Norco, Dendracin lotion, and Promolaxin. As of 6/4/2015 report, she is temporarily, totally disabled.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Follow Up Visit: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation American College of Occupational and Environmental Medicine, Chapter 7, page 127.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints.

Decision rationale: The ACOEM Chapter 12 on Low Back indicates that specialized treatments or referrals require a rationale for their use. According to the documents available for review, there is no rationale provided to support the request for an unspecified follow up visit. It is unclear from the request, which consultant the IW is to follow up with. Therefore at this time the requirements for treatment have not been met, and medical necessity has not been established.

Anaprox 550mg quantity 60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Non Steroidal Anti Inflammatory Drugs Page(s): 67-73.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS Page(s): 67, 70-73.

Decision rationale: According to the MTUS Anti-inflammatories are the traditional first line of treatment, to reduce pain so activity and functional restoration can resume, but long-term use may not be warranted. (Van Tulder-Cochrane, 2000) Recommended with cautions below. Disease-State Warnings for all NSAIDs: All NSAIDS have [U.S. Boxed Warning]: for associated risk of adverse cardiovascular events, including, MI, stroke, and new onset or worsening of pre-existing hypertension. NSAIDS should never be used right before or after a heart surgery (CABG - coronary artery bypass graft). NSAIDs can cause ulcers and bleeding in the stomach and intestines at any time during treatment (FDA Medication Guide). See NSAIDs, GI Symptoms and Cardiovascular Risks. Other disease-related concerns (non-boxed warnings): Hepatic: Use with caution in injured workers with moderate hepatic impairment and not recommended for injured workers with severe hepatic impairment. Borderline elevations of one or more liver enzymes may occur in up to 15% of injured workers taking NSAIDs. Renal: Use of NSAIDs may compromise renal function. FDA Medication Guide is provided by FDA mandate on all prescriptions dispensed for NSAIDS. Routine Suggested Monitoring: Package inserts for NSAIDs recommend periodic lab monitoring of a CBC and chemistry profile (including liver and renal function tests). There has been a recommendation to measure liver transaminases within 4 to 8 weeks after starting therapy, but the interval of repeating lab tests after this treatment duration has not been established. Routine blood pressure monitoring is recommended. Overall Dosing Recommendation: It is generally recommended that the lowest effective dose be used for all NSAIDs for the shortest duration of time consistent with the individual injured worker treatment goals. According to the documents available for review, it appears that the injured worker is taking this medication for long-term therapy of a chronic condition. Given the increased risks associated with long-term use of this medication and no documented evidence that the lowest possible dose is being used for the shortest period of time, the requirements for treatment have not been met and medical necessity has not been established.

Norco 7.5/325mg quantity 60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 80.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-97.

Decision rationale: According to the MTUS Chronic Pain Medical Treatment Guidelines section on Opioids, On-Going Management, p 74-97, (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the injured worker's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the injured worker's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain injured workers on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the injured worker should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or in injured worker treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Additionally, the MTUS states that continued use of opioids requires (a) the injured worker has returned to work, (b) the injured worker has improved functioning and pain. There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and as required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Dendracin Pain Lotion 120ml: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111-113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topicals Page(s): 111.

Decision rationale: According to the MTUS, there is little to no research to support the use of topical compounded creams. The use of these compounded agents requires knowledge of the specific analgesic effect of each agent and how it will be useful for the specific therapeutic goal required. Topical analgesics are largely experimental and there are a few randomized controlled trials to determine efficacy or safety. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

Promolaxin 11mg quantity 100: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 77.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Promolaxin.

Decision rationale: According to the ODG, Promolaxin is indicated for bowel prophylaxis of constipation secondary to chronic opioid therapy. In this case, since chronic opioid therapy is not indicated, Promolaxin is thus also not indicated. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.