

Case Number:	CM15-0089823		
Date Assigned:	05/14/2015	Date of Injury:	12/15/2008
Decision Date:	07/07/2015	UR Denial Date:	04/20/2015
Priority:	Standard	Application Received:	05/11/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Indiana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 51-year-old female, who sustained an industrial injury on 12/15/08. The injured worker has complaints of bilateral low back pain, right worse than left, right buttock pain and bilateral knee pain. The documentation noted there is tenderness to palpation over the right sacroiliac joint lower lumbar paraspinal muscles overlying the facet joints and the right greater trochanter. The diagnoses have included pain in joint, pelvic region and thigh. Treatment to date has included injections; lyrica; norco; colace; protonix; flexeril; terocin patch; effexor; wellbutrin and trazodone. The request was for hydroco/acetaminophen (Norco) 10/325 mg #180; omeprazole (Prilosec) 20 mg #60; lyrica (Pregabalin) 75 mg 2 refills; pantoprazole (Protonix) 20 mg #60 and naproxen (Anaprox) 550 mg #60.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Hydroco/APAP (Norco) 10/325 mg #180: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Shoulder, Pain, Opioids.

Decision rationale: ODG does not recommend the use of opioids for neck, low back, and shoulder pain "except for short use for severe cases, not to exceed 2 weeks." The patient has exceeded the 2 week recommended treatment length for opioid usage. MTUS does not discourage use of opioids past 2 weeks, but does state that "ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life." The treating physician does not fully document the least reported pain over the period since last assessment, intensity of pain after taking opioid, pain relief, increased level of function, or improved quality of life. Additionally, medical documents indicate that the patient has been on Norco in excess of the recommended 2-week limit. As such, the request is not medically necessary.

Omeprazole (Prilosec) 20 mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs, GI symptoms, cardiovascular risk.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs; GI Distress Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: MTUS and ODG states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily); or misoprostol (200 g four times daily) or(2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." The medical documents provided do not establish the patient as having documented GI bleeding/perforation/peptic ulcer or other GI risk factors as outlined in MTUS. As such, the request for Omeprazole is not medically necessary.

Lyrica (Pregabalin) 75 mg 2 refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines anti-epileptics.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Anti-epilepsy drugs Page(s): 16-17. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Anti-epilepsy drugs (AEDs) for pain.

Decision rationale: MTUS and ODG state, "Pregabalin (Lyrica) has been documented to be effective in treatment of diabetic neuropathy and postherpetic neuralgia, has FDA approval for both indications, and is considered first-line treatment for both. Pregabalin was also approved to treat fibromyalgia. See Anti-epilepsy drugs (AEDs) for general guidelines, as well as specific Pregabalin listing for more information and references." MTUS additionally comments "Anti-epilepsy drugs (AEDs) are also referred to as anti-convulsants. Recommended for neuropathic pain (pain due to nerve damage). A "good" response to the use of AEDs has been defined as a 50% reduction in pain and a "moderate" response as a 30% reduction. It has been reported that a 30% reduction in pain is clinically important to patients and a lack of response of this magnitude may be the "trigger" for the following: (1) a switch to a different first-line agent (TCA, SNRI or AED are considered first-line treatment); or (2) combination therapy if treatment with a single drug agent fails. (Eisenberg, 2007) (Jensen, 2006) After initiation of treatment there should be documentation of pain relief and improvement in function as well as documentation of side effects incurred with use." The patient appears to have established neuropathic pain for which Lyrica is an appropriate medication. The medical records provided do not detail any objective improvement over the last several months. Overall, pain improvement has not been documented. Given the lack of subjective and objective improvement, a request for more Lyrica is not appropriate. As such, the request is not medically necessary.

Pantoprazole (Protonix) 20 mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines PPI.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs; GI risk Page(s): 68-69. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), NSAIDs, GI symptoms & cardiovascular risk.

Decision rationale: Protonix is the brand name version of Pantoprazole, which is a proton pump inhibitor. MTUS states, "Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA)." And "Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44)." ODG states, "If a PPI is used, omeprazole OTC tablets or lansoprazole 24HR OTC are recommended for an equivalent clinical efficacy and significant cost savings. Products in this drug class have demonstrated equivalent clinical efficacy and safety at comparable doses, including esomeprazole (Nexium), lansoprazole (Prevacid), omeprazole (Prilosec), pantoprazole (Protonix), dexlansoprazole (Dexilant), and rabeprazole (Aciphex). (Shi, 2008) A trial of omeprazole or lansoprazole is recommended before Nexium therapy. The other PPIs, Protonix,

Dexilant, and Aciphex, should also be second-line. According to the latest AHRQ Comparative Effectiveness Research, all of the commercially available PPIs appeared to be similarly effective. (AHRQ, 2011)" The patient does not meet the age recommendations for increased GI risk. The medical documents provided establish the patient has experienced GI discomfort, but is nonspecific and does not indicate history of peptic ulcer, GI bleeding or perforation. Medical records do not indicate that the patient is on ASA, corticosteroids, and/or an anticoagulant; or high dose/multiple NSAID. As such, the request for Pantoprazole is not medically necessary.

Naproxen (Anaprox) 550 mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Naproxen, NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: MTUS specifies four recommendations regarding NSAID use: 1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. 2) Back Pain, Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP. 3) Back Pain, Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. 4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. Additionally, the treating physician does not document failure of primary (Tylenol) treatment. Progress notes do not indicate how long the patient has been on naproxen, but the MTUS guidelines recommend against long-term use. Dyesthesia pain is present, but as MTUS outlines, the evidence for NSAID use in neuropathic pain is inconsistent. As such, the request is not medically necessary.