

Case Number:	CM14-0119348		
Date Assigned:	09/22/2014	Date of Injury:	05/06/2002
Decision Date:	10/30/2014	UR Denial Date:	07/23/2014
Priority:	Standard	Application Received:	07/28/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Emergency Medicine and is licensed to practice in New York. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 71-year-old female who was injured on May 6, 2002. The patient continued to experience severe back pain. Physical examination was notable for decreased range of motion of lower back and altered sensory loss in the left lateral calf and bottom of her left foot. Diagnoses included low back pain and lumbosacral disc degeneration. Treatment included medications, spinal cord stimulator, psychological therapy, and surgery. Request for authorization for Norco 10/325 mg #2240 was submitted for consideration.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg #240: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Hydrocodone/Acetaminophen Chronic Pain, Opioids, Criteria for Use..

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Pain Interventions and Guidelines Page(s): 11, 74-96.

Decision rationale: Norco is the compounded medication containing Hydrocodone and Acetaminophen. Chronic Pain Medical Treatment Guidelines state that opioids are not recommended as a first line therapy. Opioid should be part of a treatment plan specific for the patient and should follow criteria for use. Criteria for use include establishment of a treatment

plan, determination if pain is nociceptive or neuropathic, failure of pain relief with non-opioid analgesics, setting of specific functional goals, and opioid contract with agreement for random drug testing. If analgesia is not obtained, opioids should be discontinued. The patient should be screened for likelihood that he or she could be weaned from the opioids if there is no improvement in pain or function. It is recommended for short term use if first-line options, such as acetaminophen or NSAIDs have failed. Opioids may be a safer choice for patients with cardiac and renal disease than antidepressants or anticonvulsants. Acetaminophen is recommended for treatment of chronic pain & acute exacerbations of chronic pain. Acetaminophen overdose is a well-known cause of acute liver failure. Hepatotoxicity from therapeutic doses is unusual. Renal insufficiency occurs in 1 to 2% of patients with overdose. The recommended dose for mild to moderate pain is 650 to 1000 mg orally every 4 hours with a maximum of 4 g/day. In this case the patient had been taking Norco since August 2012 and had not obtained analgesia. Criteria for long-term opioid use have not been met. The request should not be authorized.