

Case Number:	CM14-0120185		
Date Assigned:	09/16/2014	Date of Injury:	04/19/2011
Decision Date:	10/15/2014	UR Denial Date:	07/24/2014
Priority:	Standard	Application Received:	07/30/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Family Medicine and is licensed to practice in North Carolina. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 40-year-old with a reported date of injury of 04/19/2011. The patient has the diagnoses of chronic pain syndrome, lumbago, lumbar disc displacement L4/5 and lumbosacral neuritis. Past treatment modalities have includes physical therapy and surgical intervention in the form of a laminotomy and microdiscectomy. Per the most recent progress reports provided by the primary treating physician dated 07/15/2014, the patient had complaints of intermittent left sided low back pain that is rated a 5-7/10. The physical exam noted limited range of motion in the lumbar spine with tenderness in the paraspinal muscles. There was decreased sensation in the L5 distribution on the left and a positive straight leg raise test on the left. Treatment plan recommendations included request for a left lower extremity EMG, continuation of pain medications and a request for a psychological evaluation.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

TRAMADOL HCL 50 MG TABS: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines OPIOIDS.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids
Page(s): 78-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include:(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy.(b) The lowest possible dose should be prescribed to improve pain and function.(c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000)(d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management.(e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control.(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).(g) Continuing review of overall situation with regard to non-opioid means of pain control.(h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids(a) If the patient has returned to work(b) If the patient has improved functioning and pain(Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004)- Chronic back pain: Appears to be efficacious but limited for short-term pain relief, and long term efficacy is unclear (>16 weeks), but also appears limited. Failure to respond to a time limited. Failure to respond to a time limited course of opioids has led to the suggestion of re-assessment and consideration of alternative therapy. There is no evidence to recommend one opioid over another. In patients taking opioids for back pain, the prevalence of lifetime substance use disorders has ranged from 36% to 56% (a statistic limited by poor study design). Limited information indicated that up to one-fourth of patients who receive opioids exhibit aberrant medication-taking behavior. (Martell-Annals, 2007) (Chou, 2007) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. The most recent progress reports do note the patient has returned to work but with continued restrictions in duty. The patient continues to have significant pain without documented significant improvement in other outcome measures and function. For these reasons the criteria set forth above of ongoing and continued use of opioids have not been met. Therefore the request is not medically necessary.