

<b>Case Number:</b>	CM14-0138211		
<b>Date Assigned:</b>	09/05/2014	<b>Date of Injury:</b>	07/23/2011
<b>Decision Date:</b>	10/15/2014	<b>UR Denial Date:</b>	08/19/2014
<b>Priority:</b>	Standard	<b>Application Received:</b>	08/26/2014

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. The expert reviewer is Board Certified in Anesthesiology, has a subspecialty in Pain Medicine and is licensed to practice in Massachusetts. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/services. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

According to the documents available for review, the patient is an injured female worker. The date of injury is July 23, 2011. The patient sustained an injury to left knee. The specific mechanism of injury was not elaborated on in the notes available for review. Patient is status post left knee arthroscopic a ligament repair on March 6, 2014. The patient currently complains of pain in the bilateral knees rated 7/10 worst with mobility. The patient is maintained on the multimodal pain medication regimen including well Wellbutrin, Norco and Prozac. A request for Wellbutrin, Norco and Prozac was denied.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Retro Wellbutrin 150mg BID #60:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 27; 68; 74-96; 107.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Wellbutrin Page(s): 16.

**Decision rationale:** According to the MTUS, Bupropion (Wellbutrin), a second-generation non-tricyclic antidepressant (a noradrenaline and dopamine reuptake inhibitor) has been shown to be effective in relieving neuropathic pain of different etiologies in a small trial (41 patients).

(Finnerup, 2005) While bupropion has shown some efficacy in neuropathic pain there is no evidence of efficacy in patients with non-neuropathic chronic low back pain. (Katz, 2005) Furthermore, a recent review suggested that bupropion is generally a third-line medication for diabetic neuropathy and may be considered when patients have not had a response to a tricyclic or SNRI. (Dworkin, 2007) According to the documents available for review, the patient has none of the aforementioned indications for the use of Wellbutrin. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.

**Retro Prozac 20mg QD #30: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 27; 68; 74-96; 107.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines SSRI Page(s): 107. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG)

**Decision rationale:** According to the MTUS and the official disability guidelines, SSRIs are not effective in the treatment of chronic pain. They are effective in depression however. According to the documents available for review, the patient does not currently have a diagnosis of depression. Therefore at this time the requirements for treatment have not been met and medical necessity has not been established

**Retro Norco 10/325mg 5-6 a Day #180: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Page(s): 27; 68; 74-96; 107.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, On-Going Management Page(s): 74-97.

**Decision rationale:** should be prescribed to improve pain and function.(c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be

emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to nonopioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Additionally, the MTUS states that continued use of opioids requires (a) the patient has returned to work, (b) the patient has improved functioning and pain. There is no current documentation of baseline pain, pain score with use of opioids, functional improvement on current regimen, side effects or review of potentially aberrant drug taking behaviors as outlined in the MTUS and required for ongoing treatment. Therefore, at this time, the requirements for treatment have not been met and medical necessity has not been established.