

Case Number:	CM14-0210753		
Date Assigned:	12/23/2014	Date of Injury:	10/12/2011
Decision Date:	02/19/2015	UR Denial Date:	11/20/2014
Priority:	Standard	Application Received:	12/16/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Indiana

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This employee is a 39 year old male with date of injury of 10/12/2011 A review of the medical records indicate that the patient is undergoing treatment for discogenic cervical pain with radiculopathy and impingement of the left shoulder. Subjective complaints include 4/10 left shoulder pain with spasms, numbness, and tingling. Objective findings include limited range of motion of the cervical spine with tenderness to palpation of the paravertebrals; limited range of motion of the left shoulder with tenderness upon palpation of the rotator cuff. Treatment has included Norco, Naproxen and Flexeril. The utilization review dated 11/19/2014 non-certified Naproxen 550mg #60 and Norco 10/325 #60.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325mg quantity 60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 74-96. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG)

Neck and Upper Back (Acute and Chronic), Low Back - Lumbar & Thoracic (Acute & Chronic), Shoulder, Pain, Opioids.

Decision rationale: MTUS specifies four recommendations regarding NSAID use:1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain.2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP.3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics.4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat longterm neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. Additionally, the treating physician does not document failure of primary (Tylenol) treatment. Progress notes do not indicate how long the patient has been on Naproxen, but the MTUS guidelines recommend against long-term use. Dyesthesia pain is present, but as MTUS outlines, the evidence for NSAID use in neuropathic pain is inconsistent. As such, the request for Naproxen 550mg quantity 60 is not medically necessary.

Naproxen 550mg quantity 60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs) Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Naproxen, NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: MTUS specifies four recommendations regarding NSAID use:1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain.2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP.3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics.4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat longterm neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do

not indicate that the patient is being treated for osteoarthritis. Additionally, the treating physician does not document failure of primary (Tylenol) treatment. Progress notes do not indicate how long the patient has been on naproxen, but the MTUS guidelines recommend against long-term use. Dyesthesia pain is present, but as MTUS outlines, the evidence for NSAID use in neuropathic pain is inconsistent. As such, the request for Naproxen 550mg quantity 60 is not medically necessary.

Naproxen 550mg quantity 60; dispensed on 10/24/14: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs (non-steroidal anti-inflammatory drugs) Page(s): 67-73. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain (Chronic), Naproxen, NSAIDs (non-steroidal anti-inflammatory drugs).

Decision rationale: MTUS specifies four recommendations regarding NSAID use:1) Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain.2) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP.3) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics.4) Neuropathic pain: There is inconsistent evidence for the use of these medications to treat longterm neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. The medical documents do not indicate that the patient is being treated for osteoarthritis. Additionally, the treating physician does not document failure of primary (Tylenol) treatment. Progress notes do not indicate how long the patient has been on Naproxen, but the MTUS guidelines recommend against long-term use. Dyesthesia pain is present, but as MTUS outlines, the evidence for NSAID use in neuropathic pain is inconsistent. As such, the request for Naproxen 550mg quantity 60, dispensed on 10/24/14 is not medically necessary.