

Case Number:	CM14-0210844		
Date Assigned:	12/23/2014	Date of Injury:	02/06/2014
Decision Date:	03/05/2015	UR Denial Date:	12/07/2014
Priority:	Standard	Application Received:	12/16/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: California, District of Columbia, Maryland
 Certification(s)/Specialty: Anesthesiology, Pain Management

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

37y/o male injured worker with date of injury 2/6/14 with related neck and back pain. Per progress report dated 11/5/14, the injured worker complained of neck and upper back pain, as well as wrist pain. Physical exam revealed tenderness to palpation of the entire spine. Exam of the cervical/thoracic/lumbar spine revealed tenderness and spasm with limited range of motion and trigger points. Positive cervical compression test, positive straight leg raise bilaterally, and bilateral wrist tenderness with positive Tinel's and Phalen's were noted. Treatment to date has included physical therapy and medication management. The date of UR decision was 12/7/14.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Menthoderm Gel 240gm for 3-12 months: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Salicylate; Topical Analgesics Page(s): 105; 111-113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 60,105, 111-113.

Decision rationale: Menthoderm is methyl salicylate and menthol. Methyl salicylate may have an indication for chronic pain in this context. Per MTUS p105, "Recommended. Topical salicylate (e.g., Ben-Gay, methyl salicylate) is significantly better than placebo in chronic pain. (Mason-BMJ, 2004)." However, the CA MTUS, ODG, National Guidelines Clearinghouse, and ACOEM provide no evidence-based recommendations regarding the topical application of menthol. It is the opinion of this IMR reviewer that a lack of endorsement, a lack of mention, inherently implies a lack of recommendation, or a status equivalent to "not recommended". Since menthol is not medically indicated, then the overall product is not indicated per MTUS as outlined below. Note the statement on page 111: Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. Regarding the use of multiple medications, MTUS p60 states "Only one medication should be given at a time, and interventions that are active and passive should remain unchanged at the time of the medication change. A trial should be given for each individual medication. Analgesic medications should show effects within 1 to 3 days, and the analgesic effect of antidepressants should occur within 1 week. A record of pain and function with the medication should be recorded. (Mens, 2005) The recent AHRQ review of comparative effectiveness and safety of analgesics for osteoarthritis concluded that each of the analgesics was associated with a unique set of benefits and risks, and no currently available analgesic was identified as offering a clear overall advantage compared with the others." Therefore, it would be optimal to trial each medication individually.