

<b>Case Number:</b>	CM14-0214538		
<b>Date Assigned:</b>	01/07/2015	<b>Date of Injury:</b>	08/03/2014
<b>Decision Date:</b>	02/23/2015	<b>UR Denial Date:</b>	12/12/2014
<b>Priority:</b>	Standard	<b>Application Received:</b>	12/22/2014

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: California  
 Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The patient is a 59 year-old woman who was injured at work on 8/3/2012. The injury was primarily to her back. She is requesting review for denial of Tramadol and Mobic. Medical records corroborate ongoing care for her injuries. These records indicate that her chronic diagnoses include the following: Lumbago; Lumbosacral Neuritis; and Lumbosacral Degenerative Disc Disease. She has been treated with physical therapy and a number of different medications to include: Opioids, NSAIDs, Tricyclic Antidepressants, Muscle Relaxants and SNRIs. In the Utilization Review process MTUS/Chronic Pain Medical Treatment Guidelines were cited in the assessment of this request. The requests were non-certified for the following reasons: For the use of Tramadol, there was insufficient documentation in support of MTUS guidelines for ongoing management for patients on opioids. For the use of Mobic, non-certification was used as there is no evidence in support of long-term use of an NSAID.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Tramadol 50 MG #240 with 1 Refill: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 76-78, 880..

**Decision rationale:** The MTUS/Chronic Pain Medical Treatment Guidelines comment on the long-term use of opioids. These guidelines have established criteria of the use of opioids for the ongoing management of pain. Actions should include: prescriptions from a single practitioner and from a single pharmacy. The lowest possible dose should be prescribed to improve pain and function. There should be an ongoing review and documentation of pain relief, functional status, appropriate medication use and side effects. Pain assessment should include: current pain, the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. There should be evidence of documentation of the "4 A's for Ongoing Monitoring." These four domains include: pain relief, side effects, physical and psychological functioning, and the occurrence of any potentially aberrant drug-related behaviors. Further, there should be consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain that does not improve on opioids in 3 months. There should be consideration of an addiction medicine consult if there is evidence of substance misuse (Pages 76-78). Finally, the guidelines indicate that for chronic back pain, the long-term efficacy of opioids is unclear. Failure to respond to a time-limited course of opioids has led to the suggestion of reassessment and consideration of alternative therapy (Page 80). Based on the review of the medical records, there is insufficient documentation in support of these stated MTUS/Chronic Pain Medical Treatment Guidelines for the ongoing use of opioids. There is insufficient documentation of the "4 A's for Ongoing Monitoring." The treatment course of opioids in this patient has extended well beyond the timeframe required for a reassessment of therapy. In summary, there is insufficient documentation to support the chronic use of an opioid in this patient. Treatment with Tramadol is not considered as medically necessary.

**Mobic 15 MG #30 with 3 Refills:** Upheld

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 67-68..

**Decision rationale:** The MTUS/Chronic Pain Medical Treatment Guidelines comment on the use of NSAIDs as a treatment modality. These guidelines state the following: Specific recommendations: Osteoarthritis (including knee and hip): Recommended at the lowest dose for the shortest period in patients with moderate to severe pain. Acetaminophen may be considered for initial therapy for patients with mild to moderate pain, and in particular, for those with gastrointestinal, cardiovascular or renovascular risk factors. NSAIDs appear to be superior to

acetaminophen, particularly for patients with moderate to severe pain. There is no evidence to recommend one drug in this class over another based on efficacy. In particular, there appears to be no difference between traditional NSAIDs and COX-2 NSAIDs in terms of pain relief. The main concern of selection is based on adverse effects. COX-2 NSAIDs have fewer GI side effects at the risk of increased cardiovascular side effects, although the FDA has concluded that long-term clinical trials are best interpreted to suggest that cardiovascular risk occurs with all NSAIDs and is a class effect (with naproxyn being the safest drug). There is no evidence of long-term effectiveness for pain or function. (Chen, 2008) (Laine, 2008) Back Pain - Acute exacerbations of chronic pain: Recommended as a second-line treatment after acetaminophen. In general, there is conflicting evidence that NSAIDs are more effective than acetaminophen for acute LBP. (van Tulder, 2006) (Hancock, 2007) For patients with acute low back pain with sciatica a recent Cochrane review (including three heterogeneous randomized controlled trials) found no differences in treatment with NSAIDs vs. placebo. In patients with axial low back pain this same review found that NSAIDs were not more effective than acetaminophen for acute low-back pain, and that acetaminophen had fewer side effects. (Roelofs-Cochrane, 2008) The addition of NSAIDs or spinal manipulative therapy does not appear to increase recovery in patients with acute low back pain over that received with acetaminophen treatment and advice from their physician. (Hancock, 2007) Back Pain - Chronic low back pain: Recommended as an option for short-term symptomatic relief. A Cochrane review of the literature on drug relief for low back pain (LBP) suggested that NSAIDs were no more effective than other drugs such as acetaminophen, narcotic analgesics, and muscle relaxants. The review also found that NSAIDs had more adverse effects than placebo and acetaminophen but fewer effects than muscle relaxants and narcotic analgesics. In addition, evidence from the review suggested that no one NSAID, including COX-2 inhibitors, was clearly more effective than another. (Roelofs-Cochrane, 2008) See also Anti-inflammatory medications. Neuropathic pain: There is inconsistent evidence for the use of these medications to treat long-term neuropathic pain, but they may be useful to treat breakthrough and mixed pain conditions such as osteoarthritis (and other nociceptive pain) in with neuropathic pain. (Namaka, 2004) (Gore, 2006) See NSAIDs, GI symptoms & cardiovascular risk; NSAIDs, hypertension and renal function. Besides the above well-documented side effects of NSAIDs, there are other less well-known effects of NSAIDs, and the use of NSAIDs has been shown to possibly delay and hamper healing in all the soft tissues, including muscles, ligaments, tendons, and cartilage. (Maroon, 2006) In this case, the records indicate that NSAIDs are being used as a long-term treatment for this patient's chronic back pain. For this reason, Mobic is not considered as a medically necessary treatment.