

Case Number:	CM14-0215809		
Date Assigned:	01/05/2015	Date of Injury:	08/10/2011
Decision Date:	03/03/2015	UR Denial Date:	12/15/2014
Priority:	Standard	Application Received:	12/23/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Michigan, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The 49 year old female injured worker suffered an industrial injury on 8/10/2011. The injuries were sustained assisting in an emergency resuscitation by the injured worker in her scope of work as a nurse. The diagnoses were cervical herniated disc, lumbar herniated disc, right shoulder impingement syndrome with subsequent arthroscopic surgery and right wrist sprain. The treatment included physical therapy, medications and surgery to the shoulder on 4/24/2014. On 10/10/2014, the injured worker had a physician visit relating she got her arm caught in a gurney resulting in the decreased range of motion of the right shoulder and positive for spasm. Physical Therapy was ordered at that visit along with a urine drug analysis, topical creams and medical foods. The provider's progress note of 12/08/2014 that correlated with the request for authorization for the urine drug screen, the medical foods and the compounded creams is illegible. Prior visits do describe the injured worker to have continuing pain in the spine and right shoulder. The pain medications prescribed included gabapentin and tramadol that do have some positive effect on the pain. The UR decision on 12/15/2014 denied the above requests. The urine drug screen was denied as the rationale for the test was eligible in the 12/8/2014 progress note. The medical foods were noncertified as there were not any documented therapeutic goals. The products must be labeled for dietary management of a specific medical disorder, condition or disease for which there are distinctive nutritional requirements. The documentation did not include that information. The topical creams requested included Ketoprofen and topical muscle relaxants that are not FDA approved for use and therefore the cream that included these preparations are noncertified as the guidelines stated any preparation that included a non-

approved ingredients are noncertified. There was no clear rationale for the use of topical medications rather than the FDA approved oral forms for this patient.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Urinalysis: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter, Urine drug testing

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, steps to avoid misuse/addiction, page(Page(s): 77-78; 94..

Decision rationale: According to MTUS guidelines, urine toxicology screens is indicated to avoid misuse/addiction. “(j) Consider the use of a urine drug screen to assess for the use or the presence of illegal drugs.” In this case, there is no documentation of drug abuse or aberrant behavior. There is no documentation of drug abuse or misuse. There is no rationale provided for requesting UDS test. Therefore, Urinalysis is not medically necessary.

Theramine #90: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Medical Food

Decision rationale: According to ODG guidelines, medical food. “Recommended as indicated below. Definition: Defined in section 5(b) of the Orphan Drug Act (21 U.S.C.360ee (b) (3)) as “a food which is formulated to be consumed or administered entirely under the supervision of a physician and which is intended for the specific dietary management of a disease or condition for which distinctive nutritional requirements, based on recognized scientific principles, are established by medical evaluation.” To be considered the product must, at a minimum, meet the following criteria: (1) the product must be a food for oral or tube feeding; (2) the product must be labeled for dietary management of a specific medical disorder, disease, or condition for which there are distinctive nutritional requirements; (3) the product must be used under medical supervision. See Food labeling; Reference Daily Intakes and Daily Reference Values; Mandatory Status of Nutrition Labeling and Nutrition Content Revision proposed rule (56 FR 60366 at 60377, November 27, 1991). Medical foods are exempted from the labeling requirements for health claims and nutrient content claims under the Nutrition Labeling and Education Act of 1990 (see 21 U.S.C. 343 (q) (5) (A) (iv)). Medical foods do not have to be registered with the FDA. (CFR, 2008) Current available medical food products: Choline: Choline is a precursor of acetylcholine. There is no known medical need for choline supplementation except for the

case of long-term parenteral nutrition or for individuals with choline deficiency secondary to liver deficiency. There is inconclusive evidence that this product is indicated for an endurance aid, memory, seizures, and transient ischemic attacks. Side effects of high-dose choline include hypotension, acute GI distress, and cholinergic side effects (such as sweating and diarrhea). A fishy odor may occur with use. (AltMedDex, 2008) (Clinical Pharmacology, 2008) Glutamic Acid: This supplement is used for treatment of hypochlohydria and achlorhydria. Treatment indications include those for impaired intestinal permeability, short bowel syndrome, cancer and critical illnesses. It is generally used for digestive disorders in complementary medicine. (AltMedDex, 2008) (Lexi-Comp, 2008) 5-hydroxytryptophan: This supplement has been found to be possibly effective in treatment of anxiety disorders, fibromyalgia, obesity and sleep disorders. It has been found to be effective for depression. In alternative medicine it has been used for depression, anxiety, insomnia, obesity, aggressive behavior, eating disorders, fibromyalgia, chronic headaches and various pain disorders. It should be used with caution in individuals using SSRI antidepressants. This product has been linked to a contaminant that causes a condition called eosinophilia-myalgia syndrome. (De Benedittis, 1985) (Klarskov, 2003) (AltMedDex, 2008) (Lexi-Comp, 2008) Gamma-aminobutyric acid (GABA): This supplement is indicated for epilepsy, spasticity and tardive dyskinesia. There is no high quality peer-reviewed literature that suggests that GABA is indicated for treatment of insomnia. Adverse reactions associated with treatment include hypertension, increased heart rate and anxiety. Dose reductions are indicated for a creatinine clearance > 60 ml/min. (AltMedDex, 2008) In this low quality RCT, with no description for the actual sleep disorder, an amino acid preparation containing both GABA and 5-hydroxytryptophan reduced time to fall asleep, decreased sleep latency, increased the duration of sleep, and improved quality of sleep. (Shell, 2009) L-Serine: There is no indication in Micromedex, Clinical Pharmacology, or AltMedDex for the use of this supplement. L-Arginine: This supplement is not indicated in current references for pain or 'inflammation.' It is indicated to detoxify urine. Other indications include in use for angina, atherosclerosis, coronary artery disease, hypertension, migraines, obesity, and metabolic syndrome. (AltMedDex, 2008) (CFSAN, 2008) (Clinical Pharmacology, 2008) (Lexi-Comp, 2008) (Micromedex, 2008) Honey & cinnamon: Recommended as an option for arthritis pain. See separate listing for Honey & cinnamon. Limbrel (flavocoxid): Under study as an option for arthritis in patients at risk of adverse effects from NSAIDs, with recent evidence that Limbrel is capable of causing acute liver injury and should be used with caution. (Chalasan, 2012) See separate listing for Limbrel (flavocoxid/ arachidonic acid). See also NSAIDs, GI symptoms & cardiovascular risk; & NSAIDs, hypertension and renal function. See also Compound drugs; Co-pack drugs; Physician-dispensed drugs; Repackaged drugs. For brand names of medical foods and their respective ingredients, see Deplin (L-methylfolate); GABAdone; Sentra PM; Theramine; Trepadone; & UltraClear.”. There is no controlled studies supporting the safety and efficacy for the use of Theramine for the treatment of pain. Furthermore, there no documentation that the patient suffered from a nutrition deficit that requires the use of Theramine. Based on the above, the prescription of Theramine #90 is not medically necessary.

Sentra PM #60: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Medical Food

Decision rationale: According to ODG guidelines, medical food. “Recommended as indicated below. Definition: Defined in section 5(b) of the Orphan Drug Act (21 U.S.C.360ee (b) (3)) as “a food which is formulated to be consumed or administered entirely under the supervision of a physician and which is intended for the specific dietary management of a disease or condition for which distinctive nutritional requirements, based on recognized scientific principles, are established by medical evaluation.” To be considered the product must, at a minimum, meet the following criteria: (1) the product must be a food for oral or tube feeding; (2) the product must be labeled for dietary management of a specific medical disorder, disease, or condition for which there are distinctive nutritional requirements; (3) the product must be used under medical supervision. See Food labeling; Reference Daily Intakes and Daily Reference Values; Mandatory Status of Nutrition Labeling and Nutrition Content Revision proposed rule (56 FR 60366 at 60377, November 27, 1991). Medical foods are exempted from the labeling requirements for health claims and nutrient content claims under the Nutrition Labeling and Education Act of 1990 (see 21 U.S.C. 343 (q) (5) (A) (iv)). Medical foods do not have to be registered with the FDA. (CFR, 2008) Current available medical food products: Choline: Choline is a precursor of acetylcholine. There is no known medical need for choline supplementation except for the case of long-term parenteral nutrition or for individuals with choline deficiency secondary to liver deficiency. There is inconclusive evidence that this product is indicated for an endurance aid, memory, seizures, and transient ischemic attacks. Side effects of high-dose choline include hypotension, acute GI distress, and cholinergic side effects (such as sweating and diarrhea). A fishy odor may occur with use. (AltMedDex, 2008) (Clinical Pharmacology, 2008) Glutamic Acid: This supplement is used for treatment of hypochlorhydria and achlorhydria. Treatment indications include those for impaired intestinal permeability, short bowel syndrome, cancer and critical illnesses. It is generally used for digestive disorders in complementary medicine. (AltMedDex, 2008) (Lexi-Comp, 2008) 5-hydroxytryptophan: This supplement has been found to be possibly effective in treatment of anxiety disorders, fibromyalgia, obesity and sleep disorders. It has been found to be effective for depression. In alternative medicine it has been used for depression, anxiety, insomnia, obesity, aggressive behavior, eating disorders, fibromyalgia, chronic headaches and various pain disorders. It should be used with caution in individuals using SSRI antidepressants. This product has been linked to a contaminant that causes a condition called eosinophilia-myalgia syndrome. (De Benedittis, 1985) (Klarskov, 2003) (AltMedDex, 2008) (Lexi-Comp, 2008) Gamma-aminobutyric acid (GABA): This supplement is indicated for epilepsy, spasticity and tardive dyskinesia. There is no high quality peer-reviewed literature that suggests that GABA is indicated for treatment of insomnia. Adverse reactions associated with treatment include hypertension, increased heart rate and anxiety. Dose reductions are indicated for a creatinine clearance > 60 ml/min. (AltMedDex, 2008) In this low quality RCT, with no description for the actual sleep disorder, an amino acid preparation containing both GABA and 5-hydroxytryptophan reduced time to fall asleep, decreased sleep latency, increased the duration of sleep, and improved quality of sleep. (Shell, 2009) L-Serine: There is no indication in Micromedex, Clinical Pharmacology, or AltMedDex for the use of this supplement. L-Arginine: This supplement is not indicated in current references for pain or 'inflammation.' It is indicated to detoxify urine. Other indications include in use for angina,

atherosclerosis, coronary artery disease, hypertension, migraines, obesity, and metabolic syndrome. (AltMedDex, 2008) (CFSAN, 2008) (Clinical Pharmacology, 2008) (Lexi-Comp, 2008) (Micromedix, 2008) Honey & cinnamon: Recommended as an option for arthritis pain. See separate listing for Honey & cinnamon. Limbrel (flavocoxid): Under study as an option for arthritis in patients at risk of adverse effects from NSAIDs, with recent evidence that Limbrel is capable of causing acute liver injury and should be used with caution. (Chalasani, 2012) See separate listing for Limbrel (flavocoxid/ arachidonic acid). See also NSAIDs, GI symptoms & cardiovascular risk; & NSAIDs, hypertension and renal function. See also Compound drugs; Co-pack drugs; Physician-dispensed drugs; Repackaged drugs. For brand names of medical foods and their respective ingredients, see Deplin (L-methylfolate); GABAdone; Sentra PM; Theramine; Trepadone; & UltraClear.” There is no controlled studies supporting the safety and efficacy for the use of Sentra for the treatment of pain. Furthermore, there no documentation that the patient suffered from a nutrition deficit that requires the use of Sentra PM # 60. Based on the above, the prescription of Sentra PM # 60 is not medically necessary.

Gabadone #60: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Medical Food

Decision rationale: According to ODG guidelines, medical food. “Recommended as indicated below. Definition: Defined in section 5(b) of the Orphan Drug Act (21 U.S.C.360ee (b) (3)) as 'a food which is formulated to be consumed or administered entirely under the supervision of a physician and which is intended for the specific dietary management of a disease or condition for which distinctive nutritional requirements, based on recognized scientific principles, are established by medical evaluation.' To be considered the product must, at a minimum, meet the following criteria: (1) the product must be a food for oral or tube feeding; (2) the product must be labeled for dietary management of a specific medical disorder, disease, or condition for which there are distinctive nutritional requirements; (3) the product must be used under medical supervision. See Food labeling; Reference Daily Intakes and Daily Reference Values; Mandatory Status of Nutrition Labeling and Nutrition Content Revision proposed rule (56 FR 60366 at 60377, November 27, 1991). Medical foods are exempted from the labeling requirements for health claims and nutrient content claims under the Nutrition Labeling and Education Act of 1990 (see 21 U.S.C. 343 (q) (5) (A) (iv)). Medical foods do not have to be registered with the FDA. (CFSAN, 2008) Current available medical food products: Choline: Choline is a precursor of acetylcholine. There is no known medical need for choline supplementation except for the case of long-term parenteral nutrition or for individuals with choline deficiency secondary to liver deficiency. There is inconclusive evidence that this product is indicated for an endurance aid, memory, seizures, and transient ischemic attacks. Side effects of high-dose choline include hypotension, acute GI distress, and cholinergic side effects (such as sweating and diarrhea). A fishy odor may occur with use. (AltMedDex, 2008) (Clinical Pharmacology, 2008) Glutamic Acid: This supplement is used for treatment of hypochlohydria and achlorhydria. Treatment

indications include those for impaired intestinal permeability, short bowel syndrome, cancer and critical illnesses. It is generally used for digestive disorders in complementary medicine. (AltMedDex, 2008) (Lexi-Comp, 2008) 5-hydroxytryptophan: This supplement has been found to be possibly effective in treatment of anxiety disorders, fibromyalgia, obesity and sleep disorders. It has been found to be effective for depression. In alternative medicine it has been used for depression, anxiety, insomnia, obesity, aggressive behavior, eating disorders, fibromyalgia, chronic headaches and various pain disorders. It should be used with caution in individuals using SSRI antidepressants. This product has been linked to a contaminant that causes a condition called eosinophilia-myalgia syndrome. (De Benedittis, 1985) (Klarskov, 2003) (AltMedDex, 2008) (Lexi-Comp, 2008) Gamma-aminobutyric acid (GABA): This supplement is indicated for epilepsy, spasticity and tardive dyskinesia. There is no high quality peer-reviewed literature that suggests that GABA is indicated for treatment of insomnia. Adverse reactions associated with treatment include hypertension, increased heart rate and anxiety. Dose reductions are indicated for a creatinine clearance > 60 ml/min. (AltMedDex, 2008) In this low quality RCT, with no description for the actual sleep disorder, an amino acid preparation containing both GABA and 5-hydroxytryptophan reduced time to fall asleep, decreased sleep latency, increased the duration of sleep, and improved quality of sleep. (Shell, 2009)L-Serine: There is no indication in Micromedex, Clinical Pharmacology, or AltMedDex for the use of this supplement. L-Arginine: This supplement is not indicated in current references for pain or 'inflammation.' It is indicated to detoxify urine. Other indications include in use for angina, atherosclerosis, coronary artery disease, hypertension, migraines, obesity, and metabolic syndrome. (AltMedDex, 2008) (CFSAN, 2008) (Clinical Pharmacology, 2008) (Lexi-Comp, 2008) (Micromedex, 2008) Honey & cinnamon: Recommended as an option for arthritis pain. See separate listing for Honey & cinnamon. Limbrel (flavocoxid): Under study as an option for arthritis in patients at risk of adverse effects from NSAIDs, with recent evidence that Limbrel is capable of causing acute liver injury and should be used with caution. (Chalasani, 2012) See separate listing for Limbrel (flavocoxid/ arachidonic acid). See also NSAIDs, GI symptoms & cardiovascular risk; & NSAIDs, hypertension and renal function. See also Compound drugs; Co-pack drugs; Physician-dispensed drugs; Repackaged drugs. For brand names of medical foods and their respective ingredients, see Deplin (L-methylfolate); GABAdone; Sentra PM; Theramine; Trepadone; & UltraClear.” There is no controlled studies supporting the safety and efficacy for the use of Gabadone for the treatment of pain. Furthermore, there no documentation that the patient suffered from a nutrition deficit that requires the use of Gabadone. Based on the above, the prescription of Gabadone is not medically necessary.

Sentra AM #60: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Pain Chapter

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation Medical Food

Decision rationale: According to ODG guidelines, medical food. “Recommended as indicated below. Definition: Defined in section 5(b) of the Orphan Drug Act (21 U.s.c.360ee (b) (3)) as 'a

food which is formulated to be consumed or administered entirely under the supervision of a physician and which is intended for the specific dietary management of a disease or condition for which distinctive nutritional requirements, based on recognized scientific principles, are established by medical evaluation.' To be considered the product must, at a minimum, meet the following criteria: (1) the product must be a food for oral or tube feeding; (2) the product must be labeled for dietary management of a specific medical disorder, disease, or condition for which there are distinctive nutritional requirements; (3) the product must be used under medical supervision. See Food labeling; Reference Daily Intakes and Daily Reference Values; Mandatory Status of Nutrition Labeling and Nutrition Content Revision proposed rule (56 FR 60366 at 60377, November 27, 1991). Medical foods are exempted from the labeling requirements for health claims and nutrient content claims under the Nutrition Labeling and Education Act of 1990 (see 21 U.S.C. 343 (q) (5) (A) (iv)). Medical foods do not have to be registered with the FDA. (CFR, 2008) Current available medical food products: Choline: Choline is a precursor of acetylcholine. There is no known medical need for choline supplementation except for the case of long-term parenteral nutrition or for individuals with choline deficiency secondary to liver deficiency. There is inconclusive evidence that this product is indicated for an endurance aid, memory, seizures, and transient ischemic attacks. Side effects of high-dose choline include hypotension, acute GI distress, and cholinergic side effects (such as sweating and diarrhea). A fishy odor may occur with use. (AltMedDex, 2008) (Clinical Pharmacology, 2008) Glutamic Acid: This supplement is used for treatment of hypochlorhydria and achlorhydria. Treatment indications include those for impaired intestinal permeability, short bowel syndrome, cancer and critical illnesses. It is generally used for digestive disorders in complementary medicine. (AltMedDex, 2008) (Lexi-Comp, 2008) 5-hydroxytryptophan: This supplement has been found to be possibly effective in treatment of anxiety disorders, fibromyalgia, obesity and sleep disorders. It has been found to be effective for depression. In alternative medicine it has been used for depression, anxiety, insomnia, obesity, aggressive behavior, eating disorders, fibromyalgia, chronic headaches and various pain disorders. It should be used with caution in individuals using SSRI antidepressants. This product has been linked to a contaminant that causes a condition called eosinophilia-myalgia syndrome. (De Benedittis, 1985) (Klarskov, 2003) (AltMedDex, 2008) (Lexi-Comp, 2008) Gamma-aminobutyric acid (GABA): This supplement is indicated for epilepsy, spasticity and tardive dyskinesia. There is no high quality peer-reviewed literature that suggests that GABA is indicated for treatment of insomnia. Adverse reactions associated with treatment include hypertension, increased heart rate and anxiety. Dose reductions are indicated for a creatinine clearance > 60 ml/min. (AltMedDex, 2008) In this low quality RCT, with no description for the actual sleep disorder, an amino acid preparation containing both GABA and 5-hydroxytryptophan reduced time to fall asleep, decreased sleep latency, increased the duration of sleep, and improved quality of sleep. (Shell, 2009) L-Serine: There is no indication in Micromedex, Clinical Pharmacology, or AltMedDex for the use of this supplement. L-Arginine: This supplement is not indicated in current references for pain or 'inflammation.' It is indicated to detoxify urine. Other indications include in use for angina, atherosclerosis, coronary artery disease, hypertension, migraines, obesity, and metabolic syndrome. (AltMedDex, 2008) (CFR, 2008) (Clinical Pharmacology, 2008) (Lexi-Comp, 2008) (Micromedex, 2008) Honey & cinnamon: Recommended as an option for arthritis pain. See separate listing for Honey & cinnamon. Limbrel (flavocoxid): Under study as an option for arthritis in patients at risk of adverse effects from NSAIDs, with recent evidence that Limbrel is capable of causing acute liver injury and should be used with caution. (Chalasan, 2012) See

separate listing for Limbrel (flavocoxid/ arachidonic acid). See also NSAIDs, GI symptoms & cardiovascular risk; & NSAIDs, hypertension and renal function. See also Compound drugs; Co-pack drugs; Physician-dispensed drugs; Repackaged drugs. For brand names of medical foods and their respective ingredients, see Deplin (L-methylfolate); GABAdone; Sentra PM; Theramine; Trepadone; & UltraClear.” There is no controlled studies supporting the safety and efficacy for the use of Sentra for the treatment of pain. Furthermore, there no documentation that the patient suffered from a nutrition deficit that requires the use of Sentra PM # 60. Based on the above, the prescription of Sentra PM # 60 is not medically necessary.

Flurbiprofen/Capsaicin/Camphor 10/0.025%/2%/1% - 120 gm: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical analgesics.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics Page(s): 111.

Decision rationale: According to MTUS, in Chronic Pain Medical Treatment guidelines section Topical Analgesics (page 111), topical analgesics are largely experimental in use with few randomized controlled trials to determine efficacy or safety. Many agents are combined to other pain medications for pain control. That is limited research to support the use of many of these agents. Furthermore, according to MTUS guidelines, any compounded product that contains at least one drug or drug class that is not recommended is not recommended. There is no documentation that all component of the prescribed topical analgesic is effective for the treatment of chronic pain. Therefore, Flurbiprofen/Capsaicin/Camphor 10/0.025%/2%/1% - 120 gm is not medically necessary.