

Case Number:	CM14-0064614		
Date Assigned:	07/11/2014	Date of Injury:	08/07/2001
Decision Date:	05/27/2015	UR Denial Date:	04/07/2014
Priority:	Standard	Application Received:	05/07/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 43 year old, female who sustained a work related injury on 8/7/01. The diagnoses have included lumbar pain and degenerative joint disease bilateral knees. The treatment has included medications. In the Worker's Compensation Report to Employer dated 4/1/14, the injured worker complains of low back and bilateral knees pain. The treatment plan is refills of medications.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Cymbalta 60mg, #60 with two refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Antidepressants for chronic pain.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Cymbalta Page(s): 48-49.

Decision rationale: The California chronic pain medical treatment guidelines section on Duloxetine states: Duloxetine (Cymbalta) - Recommended as an option in first-line treatment

option in neuropathic pain. Duloxetine (Cymbalta) is a norepinephrine and serotonin reuptake inhibitor antidepressant (SNRIs). It has FDA approval for treatment of depression, generalized anxiety disorder, and for the treatment of pain related to diabetic neuropathy, with effect found to be significant by the end of week 1 effect measured as a 30% reduction in baseline pain). The starting dose is 20-60 mg/day, and no advantage has been found by increasing the dose to twice a day, except in fibromyalgia. The medication has been found to be effective for treating fibromyalgia in women with and without depression, 60 mg once or twice daily. (Arnold, 2005) The most frequent side effects include nausea, dizziness and fatigue. GI symptoms are more common early in treatment. The side effect profile of Duloxetine is thought to be less bothersome to patients than that of tricyclic anti-depressants. Note: On October 17, 2005, Eli Lilly and the U.S. Food and Drug Administration (FDA) notified healthcare professionals of revision to the PRECAUTIONS/Hepatotoxicity section of the prescribing information for Cymbalta. Post-marketing reports of hepatic injury (including hepatitis and cholestatic jaundice) suggest that patients with preexisting liver disease who take duloxetine may have an increased risk for further liver damage. The new labeling extends the Precaution against using Cymbalta in patients with substantial alcohol use to include those patients with chronic liver disease. It is recommended that Cymbalta not be administered to patients with hepatic insufficiency. See also Antidepressants for chronic pain for general guidelines, as well as specific Duloxetine listing for more information and references. On June 13, 2008, the FDA approved a new indication for duloxetine HCl delayed-release capsules (Cymbalta; Eli Lilly and Company) for the management of fibromyalgia in adults. The FDA notes that although duloxetine was effective for reducing pain in patients with and without major depressive disorder, the degree of pain relief may have been greater in those with comorbid depression. Treatment of fibromyalgia with duloxetine should be initiated at 30 mg/day for 1 week and then uptitrated to the recommended 60-mg dose. (Waknine, 2008) Note: This drug was recently included in a list of 20 medications identified by the FDA's Adverse Event Reporting System, that are under FDA investigation. (FDA, 2008) The requested medication is a first line option in the treatment of neuropathic pain per the California MTUS. The patient does not have a diagnosis of primary neuropathic pain and therefore the request is not medically necessary.

Hydrocodone, 10/325 mg, #120 with two refills: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Hydrocodone (Vicodin, Lortab).

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids
Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to

treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000)

(d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management.

(e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control.

(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).

(g) Continuing review of overall situation with regard to non-opioid means of pain control.

(h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse.

When to Continue Opioids;

(a) If the patient has returned to work.

(b) If the patient has improved functioning and pain (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox- AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004)

The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore criteria for the ongoing use of opioids have not been met and the request is not medically necessary.