

Case Number:	CM14-0075072		
Date Assigned:	07/16/2014	Date of Injury:	06/09/2011
Decision Date:	05/27/2015	UR Denial Date:	05/21/2014
Priority:	Standard	Application Received:	05/22/2014

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 33 year old male, who sustained an industrial injury on 6/09/2011, while employed as a chef. He reported twisting his right knee and falling to the floor and hitting his left shoulder blade. The injured worker was diagnosed as having thoracic sprain/strain, knee tendinitis/bursitis, and right medial meniscal tear. Treatment to date has included diagnostics, right knee surgery x2 (2012), physical therapy, cortisone injections, and medications. In 10/2013, current medications included Celebrex, Tramadol, Norco, and Robaxin. Urine drug screen, dated 12/12/2013, was inconsistent with expected results. On 5/01/2014, the injured worker complained of pain in the thoracic area and bilateral knees (not rated). Physical exam noted decreased range of motion over the thoracic spine, with spasm, tenderness, and guarding. Tenderness to palpation was noted over the left knee and right knee medial joint lines. McMurray's tests were positive. The treatment plan included Neurontin, Terocin patch, and Norflex. His work status was permanent and stationary.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

One prescription for Neurontin 300mg, #90 with one refill: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Criteria /Guidelines Gabapentin.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines gabapentin Page(s): 18.

Decision rationale: The California chronic pain medical treatment guidelines section on Neurontin states: Gabapentin (Neurontin, Gabarone, generic available) has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain. (Backonja, 2002) (ICSI, 2007) (Knotkova, 2007) (Eisenberg, 2007) (Attal, 2006) This RCT concluded that gabapentin monotherapy appears to be efficacious for the treatment of pain and sleep interference associated with diabetic peripheral neuropathy and exhibits positive effects on mood and quality of life. (Backonja, 1998) It has been given FDA approval for treatment of post-herpetic neuralgia. The number needed to treat (NNT) for overall neuropathic pain is 4. It has a more favorable side-effect profile than Carbamazepine, with a number needed to harm of 2.5. (Wiffen2-Cochrane, 2005) (Zaremba, 2006) Gabapentin in combination with morphine has been studied for treatment of diabetic neuropathy and postherpetic neuralgia. When used in combination the maximum tolerated dosage of both drugs was lower than when each was used as a single agent and better analgesia occurred at lower doses of each. (Gilron-NEJM, 2005) Recommendations involving combination therapy require further study. The requested medication is a first line agent to treatment neuropathic pain. The patient does not have a diagnosis of neuropathic pain. Therefore the request is not medically necessary.

One prescription for Norflex 100mg, #60 with one refill: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines muscle relaxants Page(s): 63-65.

Decision rationale: The California chronic pain medical treatment guidelines section on muscle relaxants states: Recommend non-sedating muscle relaxants with caution as a second-line option for short-term treatment of acute exacerbations in patients with chronic LBP. (Chou, 2007) (Mens, 2005) (Van Tulder, 1998) (van Tulder, 2003) (van Tulder, 2006) (Schnitzer, 2004) (See, 2008) Muscle relaxants may be effective in reducing pain and muscle tension, and increasing mobility. However, in most LBP cases, they show no benefit beyond NSAIDs in pain and overall improvement. Also there is no additional benefit shown in combination with NSAIDs. Efficacy appears to diminish over time, and prolonged use of some medications in this class may lead to dependence. (Homik, 2004) (Chou, 2004) This medication is not intended for long-term use per the California MTUS. The medication has not been prescribed for the flare-up of chronic low back pain. This is not an approved use for the medication. For these reasons, criteria for the use of this medication have not been met. Therefore the request is not medically necessary.

One prescription for Terocin Patches, #30 with one refill: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Topical Analgesics.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines topical analgesics Page(s): 111-113.

Decision rationale: The California chronic pain medical treatment guidelines section on topical analgesics states: Recommended as an option as indicated below. Largely experimental in use with few randomized controlled trials to determine efficacy or safety. Primarily recommended for neuropathic pain when trials of antidepressants and anticonvulsants have failed. (Namaka, 2004) These agents are applied locally to painful areas with advantages that include lack of systemic side effects, absence of drug interactions, and no need to titrate. (Colombo, 2006) Many agents are compounded as monotherapy or in combination for pain control (including NSAIDs, opioids, capsaicin, local anesthetics, antidepressants, glutamate receptor antagonists, -adrenergic receptor agonist, adenosine, cannabinoids, cholinergic receptor agonists, agonists, prostanoids, bradykinin, adenosine triphosphate, biogenic amines, and nerve growth factor). (Argoff, 2006) There is little to no research to support the use of many of these agents. Any compounded product that contains at least one drug (or drug class) that is not recommended is not recommended. The requested medication contains ingredients, which are not indicated per the California MTUS for topical analgesic use. Therefore the request is not medically necessary.