

Case Number:	CM15-0004025		
Date Assigned:	01/15/2015	Date of Injury:	08/07/2008
Decision Date:	03/10/2015	UR Denial Date:	12/31/2014
Priority:	Standard	Application Received:	01/08/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 60 year old male who sustained an industrial injury on August 7, 2008. He reports pain to the right leg, bilateral low back, and right foot/ankle and has been diagnosed with post laminectomy syndrome lumbar region and back pain, lumbar, with radiculopathy. Treatment to date has included epidural steroid injection, pain medications, cognitive behavioral psychotherapy, acupuncture, and physical therapy. Currently the injured worker complains of pain that is worse with lifting, stress, twisting, weather changes, and cold. The treatment plan included medications and exercise. On December 31, 2014 Utilization Review modified percocet 10/325 mg # 120 with 1 refill and non certified 1 toradol injection citing the MTUS guidelines.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Percocet 10/325mg #120: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids
Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include:(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy.(b) The lowest possible dose should be prescribed to improve pain and function.(c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or nonadherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000)(d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management.(e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control.(f) Documentation of misuse of medications (doctor-shopping, uncontrolled drug escalation, drug diversion).(g) Continuing review of overall situation with regard to nonopioid means of pain control.(h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids(a) If the patient has returned to work(b) If the patient has improved functioning and pain(Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. The most recent documentation provided from the primary treating physician has no evidence of subjective improvement in pain such as VAS scores. There is also no objective measure of improvement in function. The patient was switched to Percocet from oxycontin for decreased reliance on opioids but the criteria for ongoing opioid therapy have not been met. Therefore the request is not certified.

Toradol injection: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDS
Page(s): 72. Decision based on Non-MTUS Citation Toradol

Decision rationale: The California chronic pain medical treatment guidelines section on Ketorolac states: Ketorolac (Toradol, generic available): 10 mg. [Boxed Warning]: This medication is not indicated for minor or chronic painful conditions. Per the ODG: Only recommended for short-term in management of moderately severe acute pain that requires analgesia at the opioid level. In this case, the documentation does not indicate acute pain treatment but rather than the treatment of a chronic pain condition. In the absence of acute pain treatment, the medication is not indicated per the California MTUS and the ODG. Therefore the request is not certified.