

Case Number:	CM15-0100172		
Date Assigned:	06/02/2015	Date of Injury:	09/23/2013
Decision Date:	06/30/2015	UR Denial Date:	05/12/2015
Priority:	Standard	Application Received:	05/26/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 67 year old, male who sustained a work related injury on 9/23/13. He was going out a swinging door and someone was coming in the door with a two-wheeled cart full of wine and slammed into the injured worker. The diagnoses have included left wrist arthritis, scapholunate ligament tear, possible left cubital tunnel syndrome and triangular fibrocartilage complex tear. Treatments not noted. In the PR-2 dated 4/28/15, the injured worker complains of left wrist pain, weakness, numbness and tingling into the ring and small fingers. He has tenderness over the left radiocarpal joint dorsally. The left index through small fingers have ulnar drift at the metacarpophalangeal joint, approximately 10-20 degrees. The treatment plan includes a request for an EMG/NCV testing.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Electromyogram (EMG) of left median ulnar wrist: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 10 Elbow Disorders (Revised 2007).

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 173-174.

Decision rationale: The ACOEM chapter on neck and upper back complaints and special diagnostic studies states: Criteria for ordering imaging studies are: Emergence of a red flag, Physiologic evidence of tissue insult or neurologic dysfunction, Failure to progress in a strengthening program intended to avoid surgery, Clarification of the anatomy prior to an invasive procedure. Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The assessment may include sensory-evoked potentials (SEPs) if spinal stenosis or spinal cord myelopathy is suspected. If physiologic evidence indicates tissue insult or nerve impairment, consider a discussion with a consultant regarding next steps, including the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computer tomography [CT] for bony structures). Additional studies may be considered to further define problem areas. The recent evidence indicates cervical disk annular tears may be missed on MRIs. The clinical significance of such a finding is unclear, as it may not correlate temporally or anatomically with symptoms. The provided clinical documentation for review meets criteria as cited above for the requested service and therefore it is medically necessary.

Nerve Conduction Velocity (NCV) of left median ulnar wrist: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 10 Elbow Disorders (Revised 2007).

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 173-174.

Decision rationale: The ACOEM chapter on neck and upper back complaints and special diagnostic studies states: Criteria for ordering imaging studies are:-Emergence of a red flag- Physiologic evidence of tissue insult or neurologic dysfunction-Failure to progress in a strengthening program intended to avoid surgery-Clarification of the anatomy prior to an invasive procedure Physiologic evidence may be in the form of definitive neurologic findings on physical examination, electrodiagnostic studies, laboratory tests, or bone scans. Unequivocal findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging studies if symptoms persist. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction can be obtained before ordering an imaging study. Electromyography (EMG), and nerve conduction velocities (NCV), including H-reflex tests, may help identify subtle focal neurologic dysfunction in patients with neck or arm symptoms, or both, lasting more than three or four weeks. The assessment may include sensory-evoked potentials (SEPs) if spinal stenosis or spinal cord myelopathy is

suspected. If physiologic evidence indicates tissue insult or nerve impairment, consider a discussion with a consultant regarding next steps, including the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computer tomography [CT] for bony structures). Additional studies may be considered to further define problem areas. The recent evidence indicates cervical disk annular tears may be missed on MRIs. The clinical significance of such a finding is unclear, as it may not correlate temporally or anatomically with symptoms. The provided clinical documentation for review meets criteria as cited above for the requested service and therefore it is medically necessary.