

Case Number:	CM15-0104735		
Date Assigned:	06/09/2015	Date of Injury:	03/24/2015
Decision Date:	07/15/2015	UR Denial Date:	05/08/2015
Priority:	Standard	Application Received:	06/01/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 29-year-old male, who sustained an industrial injury on March 24, 2015, incurring back, left shoulder and arm injuries. He was diagnosed with thoracic sprain, lumbar sprain, bilateral shoulder sprain, left elbow epicondylitis, bilateral wrist sprain and bilateral knee sprain. Treatment included anti-inflammatory drugs, anti-anxiety medications, sleep aides, proton pump inhibitor, and physical therapy and chiropractic sessions. Currently, the injured worker complained of persistent chronic upper and lower back pain, shoulder pain, elbow and wrist pain, sleep interruption, depression and anxiety. The treatment plan that was requested for authorization included a prescription for Omeprazole.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Omeprazole 20mg, #30: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 10 Elbow Disorders (Revised 2007), Chapter 11 Forearm, Wrist, and Hand Complaints, Chapter 3 Initial Approaches to Treatment, Chapter 12 Low Back Complaints, Chapter 13 Knee Complaints Page(s): 34, 601, 265, 268, 301, 303, 341, Chronic Pain Treatment Guidelines NSAIDs, GI symptoms & cardiovascular risk; Topical Analgesics Page(s): 68, 111-113.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines
NSAID Page(s): 68-72.

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID therapy and proton pump inhibitors (PPI) states: Recommend with precautions as indicated below. Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastro duodenal lesions. Recommendations Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200g four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if necessary. There is no documentation provided that places this patient at intermediate or high risk that would justify the use of a PPI. There is no mention of current gastrointestinal or cardiovascular disease. For these reasons, the criteria set forth above per the California MTUS for the use of this medication has not been met. Therefore, the request is not medically necessary.