

<b>Case Number:</b>	CM15-0110466		
<b>Date Assigned:</b>	06/17/2015	<b>Date of Injury:</b>	05/10/2012
<b>Decision Date:</b>	07/15/2015	<b>UR Denial Date:</b>	05/28/2015
<b>Priority:</b>	Standard	<b>Application Received:</b>	06/08/2015

### HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:  
 State(s) of Licensure: North Carolina  
 Certification(s)/Specialty: Family Practice

### CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 35 year old male sustained an industrial injury to the left leg and both wrists and hands on 5/10/12. Previous treatment included bilateral carpal tunnel release, electromyography, physical therapy, splinting, occupational therapy and medications. In the most recent documentation submitted for review, an Agreed Medical Evaluation dated 3/24/15, the injured worker complained of bilateral wrist, hand and thumb pain with radiation to the fingers associated with weakness and left lower leg tenderness and sensitivity. Physical exam was remarkable for tenderness to palpation to bilateral wrists and hands with decreased sensation to light touch in the lateral left forearm and right thumb, index and ring fingers with positive right Tinel's and Phalen's and positive left Tinel's and questionable positive left Phalen's. Current medications included Celebrex, Gabapentin, Elavil and Voltaren gel. Current diagnoses included possible left elbow ulnar neuritis, right carpal tunnel release and de Quervain's release with residual carpal tunnel syndrome, left carpal tunnel release with residual carpal tunnel syndrome, left middle trigger finger and left lateral leg foreign body removal. The physician recommended work modification, short courses of physical therapy, local injections, splinting and ongoing medication management.

### IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

**Celebrex 100mg #60: Upheld**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 67-68.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines NSAID Page(s): 68-72.

**Decision rationale:** The California chronic pain medical treatment guidelines section on NSAID use and proton pump inhibitors (PPI) states: Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastroduodenal lesions. Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g., ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 g four times daily); or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox- 2 selective agent plus a PPI if absolutely necessary. Cardiovascular disease: A non- pharmacological choice should be the first option in patients with cardiac risk factors. It is then suggested that acetaminophen or aspirin be used for short term needs. An opioid also remains a short-term alternative for analgesia. Major risk factors (recent MI, or coronary artery surgery, including recent stent placement): If NSAID therapy is necessary, the suggested treatment is naproxen plus low-dose aspirin plus a PPI. Mild to moderate risk factors: If long-term or high- dose therapy is required, full-dose naproxen (500 mg twice a day) appears to be the preferred choice of NSAID. If naproxen is ineffective, the suggested treatment is: (1) the addition of aspirin to naproxen plus a PPI, or (2) a low-dose Cox-2 plus ASA. Cardiovascular risk does appear to extend to all non-aspirin NSAIDs, with the highest risk found for the Cox-2 agents. (Johnsen, 2005) (Lanas, 2006) (Antman, 2007) (Laine, 2007) Use with Aspirin for cardio protective effect: In terms of GI protective effect: The GI protective effect of Cox-2 agents is diminished in patients taking low-dose aspirin and a PPI may be required for those patients with GI risk factors. (Laine, 2007) In terms of the actual cardio protective effect of aspirin: Traditional NSAIDs (both ibuprofen and naproxen) appear to attenuate the antiplatelet effect of enteric- coated aspirin and should be taken 30 minutes after ASA or 8 hours before. (Antman, 2007) Cox-2 NSAIDs and diclofenac (a traditional NSAID) do not decrease anti-platelet effect. (Laine, 2007) The patient does not have risk factors that would require a COX-2 inhibitor over a traditional NSAID. Therefore the request is not medically necessary.

**Neurontin 600mg #690: Overturned**

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Anti-epilepsy drugs.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines gabapentin Page(s): 18.

**Decision rationale:** The California chronic pain medical treatment guidelines section on Neurontin states: Gabapentin (Neurontin, Gabarone, generic available) has been shown to be effective for treatment of diabetic painful neuropathy and postherpetic neuralgia and has been considered as a first-line treatment for neuropathic pain. (Backonja, 2002) (ICSI, 2007) (Knotkova, 2007) (Eisenberg, 2007) (Attal, 2006) This RCT concluded that gabapentin monotherapy appears to be efficacious for the treatment of pain and sleep interference associated with diabetic peripheral neuropathy and exhibits positive effects on mood and quality of life. (Backonja, 1998) It has been given FDA approval for treatment of post-herpetic neuralgia. The number needed to treat (NNT) for overall neuropathic pain is 4. It has a more favorable side-effect profile than Carbamazepine, with a number needed to harm of 2.5. (Wiffen2-Cochrane, 2005) (Zaremba, 2006) Gabapentin in combination with morphine has been studied for treatment of diabetic neuropathy and postherpetic neuralgia. When used in combination the maximum tolerated dosage of both drugs was lower than when each was used as a single agent and better analgesia occurred at lower doses of each. (Gilron-NEJM, 2005) Recommendations involving combination therapy require further study. The patient has the diagnosis of neuropathic pain. Therefore the request is necessary and approved.

**Elavil 25mg #30:** Overturned

**Claims Administrator guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines Antidepressants for chronic pain Page(s): 13-16.

**MAXIMUS guideline:** Decision based on MTUS Chronic Pain Treatment Guidelines antidepressants Page(s): 15.

**Decision rationale:** The California MTUS section on antidepressants states: Tricyclic antidepressants are recommended over selective serotonin reuptake inhibitors (SSRIs), unless adverse reactions are a problem. Caution is required because tricyclics have a low threshold for toxicity, and tricyclic antidepressant overdose is a significant cause of fatal drug poisoning due to their cardiovascular and neurological effects. Tricyclic antidepressants have been shown in both a meta-analysis (McQuay, 1996) and a systematic review (Collins, 2000) to be effective, and are considered a first-line treatment for neuropathic pain. (Namaka, 2004) (Dworkin, 2003) (Gilron, 2006) (Wolfe, 2004) (Dworkin, 2007) (Saarto-Cochrane, 2007) This class of medications works in both patients with normal mood and patients with depressed mood when used in treatment for neuropathic pain. (Sindrup, 2005) Indications in controlled trials have shown effectiveness in treating central post-stroke pain, post-herpetic neuralgia (Argoff, 2004), painful diabetic and non-diabetic polyneuropathy, and post-mastectomy pain. Negative results were found for spinal cord pain and phantom-limb pain, but this may have been due to study design. (Finnerup, 2005) Tricyclics have not demonstrated significance in randomized-control trials in treating HIV neuropathy, spinal cord injury, cisplatin neuropathy, neuropathic cancer pain, phantom limb pain or chronic lumbar root pain. (Dworkin, 2007) One review reported the NNT for at least moderate neuropathic pain relief with tricyclics is 3.6 (3-4.5), with the NNT for amitriptyline being 3.1 (2.5-4.2). The NNT for venlafaxine calculated using 3

studies was reported to be 3.1 (2.2-5.1). (Saarto-Cochrane, 2007) Another review reported that the NNT for 50% improvement in neuropathic pain was 2 to 3 for tricyclic antidepressants, 4 for venlafaxine, and 7 for SSRIs (Perrot, 2008). The patient has neuropathic pain and therefore the request is medically necessary as this is a first line treatment option.