

Case Number:	CM15-0110554		
Date Assigned:	06/17/2015	Date of Injury:	08/07/2013
Decision Date:	07/15/2015	UR Denial Date:	05/28/2015
Priority:	Standard	Application Received:	06/08/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: North Carolina
 Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 46-year-old male, who sustained an industrial injury on 8/7/13. He reported right shoulder and elbow injury following from a ladder. The injured worker was diagnosed as having status post right shoulder surgery with persistent pain and complaint of right elbow pain. Treatment to date has included right shoulder surgery, chiropractic treatment, physical therapy, oral medications including Celebrex, activity restrictions and home exercise program. Currently, the injured worker complains of constant right elbow pain rated 6/10 associated with numbness and tingling in the forearm and constant right shoulder pain rated 5/10, worse with reaching, lifting, pushing and pulling. He is temporarily totally disabled. Physical exam noted well healed incision of right shoulder with restricted range of motion and tenderness anteriorly and laterally and right elbow tenderness medially and laterally at the epicondyle with positive Cozen's. A request for authorization was submitted for Celebrex 200mg #30, Ambien 10mg #50 and ultrasound of right arm to rule out lipoma.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Celebrex 200mg #30 x 1 refill: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAIDs Page(s): 63.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines NSAID Page(s): 68-72.

Decision rationale: The California chronic pain medical treatment guidelines section on NSAID use and proton pump inhibitors (PPI) states: Clinicians should weight the indications for NSAIDs against both GI and cardiovascular risk factors. Determine if the patient is at risk for gastrointestinal events: (1) age > 65 years; (2) history of peptic ulcer, GI bleeding or perforation; (3) concurrent use of ASA, corticosteroids, and/or an anticoagulant; or (4) high dose/multiple NSAID (e.g., NSAID + low-dose ASA). Recent studies tend to show that H. Pylori does not act synergistically with NSAIDS to develop gastroduodenal lesions. Recommendations: Patients with no risk factor and no cardiovascular disease: Non-selective NSAIDs OK (e.g, ibuprofen, naproxen, etc.) Patients at intermediate risk for gastrointestinal events and no cardiovascular disease: (1) A non-selective NSAID with either a PPI (Proton Pump Inhibitor, for example, 20 mg omeprazole daily) or misoprostol (200 mg four times daily) or (2) a Cox-2 selective agent. Long-term PPI use (> 1 year) has been shown to increase the risk of hip fracture (adjusted odds ratio 1.44). Patients at high risk for gastrointestinal events with no cardiovascular disease: A Cox-2 selective agent plus a PPI if absolutely necessary. Cardiovascular disease: A non-pharmacological choice should be the first option in patients with cardiac risk factors. It is then suggested that acetaminophen or aspirin be used for short-term needs. An opioid also remains a short-term alternative for analgesia. Major risk factors (recent MI, or coronary artery surgery, including recent stent placement): If NSAID therapy is necessary, the suggested treatment is naproxyn plus low-dose aspirin plus a PPI. Mild to moderate risk factors: If long-term or high- dose therapy is required, full-dose naproxen (500 mg twice a day) appears to be the preferred choice of NSAID. If naproxyn is ineffective, the suggested treatment is: (1) the addition of aspirin to naproxyn plus a PPI, or (2) a low-dose Cox-2 plus ASA. Cardiovascular risk does appear to extend to all non-aspirin NSAIDs, with the highest risk found for the Cox-2 agents. (Johnsen, 2005) (Lanas, 2006) (Antman, 2007) (Laine, 2007) Use with Aspirin for cardioprotective effect: In terms of GI protective effect: The GI protective effect of Cox-2 agents is diminished in patients taking low-dose aspirin and a PPI may be required for those patients with GI risk factors. (Laine, 2007) In terms of the actual cardio protective effect of aspirin: Traditional NSAIDs (both ibuprofen and naproxen) appear to attenuate the antiplatelet effect of enteric-coated aspirin and should be taken 30 minutes after ASA or 8 hours before. (Antman, 2007) Cox-2 NSAIDs and diclofenac (a traditional NSAID) do not decrease anti-platelet effect. (Laine, 2007) The patient does not have risk factors that would require a COX-2 inhibitor over a traditional NSAID. Therefore, the request is not medically necessary.

Ambien 10mg #50: Upheld

Claims Administrator guideline: The Claims Administrator did not base their decision on the MTUS. Decision based on Non-MTUS Citation Official Disability Guidelines, Pain, Insomnia.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation ODG, ambien.

Decision rationale: The California MTUS and the ACOEM do not specifically address the requested medication. Per the ODG: Zolpidem is a prescription short-acting non-benzodiazepine hypnotic approved for the short-term treatment of insomnia. Proper sleep hygiene is critical to the individual with chronic pain. While sleeping pills, so-called minor tranquilizers and anti-anxiety medications are commonly prescribed in chronic pain, pain specialists rarely, if ever, recommend them for long-term use. There is also concern that they may increase pain and depression over the long-term. The medication is not intended for use greater than 6 weeks. There is no notation or rationale given for longer use in the provided progress reports. There is no documentation of other preferred long-term insomnia intervention choices being tried and failed. For these reasons, the request is not medically necessary.

Ultrasound of right arm to rule out lipoma: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: The Expert Reviewer did not base their decision on the MTUS. Decision based on Non-MTUS Citation up-to date guidelines, lipoma.

Decision rationale: The California MTUS, ODG and the ACOEM do not specifically address the requested service. The up-to date clinical guidelines section on lipomas indicates that most lipomas do not require any imaging studies unless there is clinical suspicion by exam of the lipoma compromising other surrounding tissue. This criterion has not been met in the provided clinical documentation for review and therefore the request is not medically necessary.