

Case Number:	CM15-0110708		
Date Assigned:	06/17/2015	Date of Injury:	07/16/2013
Decision Date:	07/15/2015	UR Denial Date:	05/29/2015
Priority:	Standard	Application Received:	06/09/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 48 year old female sustained an industrial injury to the low back. Documentation did not disclose previous treatments or the date of injury. In the only documentation submitted for review, a PR-2 dated 5/21/15, the injured worker complained of low back pain with numbness down the right leg. Physical exam was remarkable for lumbar spine with tenderness to palpation, muscle spasms, decreased range of motion and decreased sensation to the right L5-S1 distribution. The physician note that x-rays of the lower extremities were within normal limits. Magnetic resonance imaging of the lumbar spine was negative for herniated nucleus pulposus. Current diagnosis included lumbar spine sprain/strain. The treatment plan included requesting authorization for physical therapy and right leg electromyography and returning to work on modified duty on 5/21/15.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Outpatient physical therapy to lumbar 3 times a week for 4 weeks: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Physical Medicine.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines physical medicine Page(s): 98-99.

Decision rationale: The California chronic pain medical treatment guidelines section on physical medicine states: Recommended as indicated below. Passive therapy (those treatment modalities that do not require energy expenditure on the part of the patient) can provide short-term relief during the early phases of pain treatment and are directed at controlling symptoms such as pain, inflammation and swelling and to improve the rate of healing soft tissue injuries. They can be used sparingly with active therapies to help control swelling, pain and inflammation during the rehabilitation process. Active therapy is based on the philosophy that therapeutic exercise and/or activity are beneficial for restoring flexibility, strength, endurance, function, range of motion, and can alleviate discomfort. Active therapy requires an internal effort by the individual to complete a specific exercise or task. This form of therapy may require supervision from a therapist or medical provider such as verbal, visual and/or tactile instruction(s). Patients are instructed and expected to continue active therapies at home as an extension of the treatment process in order to maintain improvement levels. Home exercise can include exercise with or without mechanical assistance or resistance and functional activities with assistive devices. (Colorado, 2002) (Airaksinen, 2006) Patient-specific hand therapy is very important in reducing swelling, decreasing pain, and improving range of motion in CRPS. (Li, 2005) The use of active treatment modalities (e.g., exercise, education, activity modification) instead of passive treatments is associated with substantially better clinical outcomes. In a large case series of patients with low back pain treated by physical therapists, those adhering to guidelines for active rather than passive treatments incurred fewer treatment visits, cost less, and had less pain and less disability. The overall success rates were 64.7% among those adhering to the active treatment recommendations versus 36.5% for passive treatment. (Fritz, 2007) Physical Medicine Guidelines, Allow for fading of treatment frequency (from up to 3 visits per week to 1 or less), plus active self-directed home Physical Medicine. Myalgia and myositis, unspecified (ICD9 729.1): 9-10 visits over 8 weeks Neuralgia, neuritis, and radiculitis, unspecified (ICD9 729.2) 8-10 visits over 4 weeks Reflex sympathetic dystrophy (CRPS) (ICD9 337.2): 24 visits over 16 weeks. The requested amount of physical therapy is in excess of California chronic pain medical treatment guidelines. There is no objective explanation why the patient would need excess physical therapy and not be transitioned to active self-directed physical medicine. The request is not medically necessary.

EMG/NCV bilateral lower extremities to rule out radiculopathy: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG), Low Back Chapter, NCS.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 12 Low Back Complaints Page(s): 303.

Decision rationale: The ACOEM chapters on low back complaints and the need for lower extremity EMG/NCV states: Unequivocal objective findings that identify specific nerve compromise on the neurologic examination are sufficient evidence to warrant imaging in patients

who do not respond to treatment and who would consider surgery an option. When the neurologic examination is less clear, however, further physiologic evidence of nerve dysfunction should be obtained before ordering an imaging study. Indiscriminant imaging will result in false- positive findings, such as disk bulges, that are not the source of painful symptoms and do not warrant surgery. If physiologic evidence indicates tissue insult or nerve impairment, the practitioner can discuss with a consultant the selection of an imaging test to define a potential cause (magnetic resonance imaging [MRI] for neural or other soft tissue, computer tomography [CT] for bony structures). Electromyography (EMG), including H-reflex tests, may be useful to identify subtle, focal neurologic dysfunction in patients with low back symptoms lasting more than three or four weeks. There are unequivocal objective findings of nerve compromise on the neurologic exam provided for review. However, there is not mention of surgical consideration. There are no unclear neurologic findings on exam. For these reasons, criteria for lower extremity EMG/NCV have not been met as set forth in the ACOEM. Therefore, the request is not medically necessary.