

Case Number:	CM15-0110773		
Date Assigned:	06/18/2015	Date of Injury:	05/19/2013
Decision Date:	07/16/2015	UR Denial Date:	05/12/2015
Priority:	Standard	Application Received:	06/09/2015

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:
 State(s) of Licensure: New Jersey, Alabama, California
 Certification(s)/Specialty: Neurology, Neuromuscular Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

This 58 year old woman sustained an industrial injury on 5/19/2013. The mechanism of injury is not detailed. Diagnoses include status post right hip surgery, suspect sacroiliac joint instability, and progressive lumbar degenerative disc disease. Treatment has included oral medications. Physician notes dated 4/28/2015 show complaints of continued low back pain with radiation to the ischium and groin and increased popping and snapping. Recommendations include physiatry consultation, second opinion orthopedic consultation, continue current medication regimen, and follow up in six weeks.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Oxycodone 5mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids, Criteria for Use.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Opioids Page(s): 75-81.

Decision rationale: According to MTUS guidelines, Oxycodone as well as other short acting opioids are indicated for intermittent or breakthrough pain. It can be used in acute post operative pain. It is not recommended for chronic pain of long term use as prescribed in this case. In addition and according to MTUS guidelines, ongoing use of opioids should follow specific rules: "(a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework." There is no documentation that the patient has pain breakthrough. There is no documentation of pain and functional improvement with previous use of opioids. There is no rationale for a continuous use of Oxycodone. Therefore, the prescription of Oxycodone 5mg #60 is not medically necessary.

Diazepam 5mg #60: Upheld

Claims Administrator guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Muscle Relaxants (for pain) Page(s): 63-66.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines Benzodiazepines Page(s): 24.

Decision rationale: According to MTUS guidelines, benzodiazepines are not recommended for long term use for pain management because of unproven long term efficacy and because of the risk of dependence. Most guidelines limit their use to 4 weeks. There is no recent documentation that the patient has insomnia and that Diazepam was effective in the past. Therefore, the prescription of Valium (Diazepam) 5mg #60 is not medically necessary.