

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM15-0111962 | | |
| Date Assigned: | 06/19/2015 | Date of Injury: | 05/29/2012 |
| Decision Date: | 07/17/2015 | UR Denial Date: | 05/22/2015 |
| Priority: | Standard | Application Received: | 06/10/2015 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: North Carolina

Certification(s)/Specialty: Family Practice

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 56 year old male patient who sustained an industrial injury on 05/29/2012. On 12/01/2014 the patient underwent a magnetic resonance imaging study of the cervical spine that revealed multi-level degenerative changes of the cervical spine with multi-level spinal canal and neural foraminal compromise. Previous treatment modality to include: physical therapy, medication management, extracorporeal shockwave therapy. A secondary treating office visit dated 12/10/2014 reported the patient with subjective complaint of symptom being unchanged. The objective assessment found the lumbar spine with pain to palpation and decreased range of motion. The following diagnoses were applied: displacement cervical disc without myelopathy; strain/sprain of shoulder/arm, and lumbar intervertebral disc syndrome. The plan of care involved: continuing with conservative treatment as the patient is not interested in surgical intervention at this time. A primary treating office visit dated 11/18/2014 reported the patient suffering from chronic pain in his cervical spine and right shoulder. There is recommendation for a neurological consultation along with a surgical consultation regarding hernia. Current subjective complaints consist of: constant severe cervical spine pain radiating into bilateral shoulders; daily headaches located over bilateral parietal lobes accompanied with blurred vision; right constant shoulder pain; constant low back pain radiating into bilateral hips; difficulty sleeping; stress, anxiety, depression, inguinal pain, skin irritation, and hearing loss. The following diagnoses were applied: cervical spine myoligamentous injury; posttraumatic headaches rule out traumatic brain injury; right shoulder myoligamentous injury; lumbar spine herniated nucleus pulposus; secondary difficulty sleeping, and secondary stress, anxiety and depression. The patient is temporarily totally disabled.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

Norco 10/325 mg #90: Upheld

Claims Administrator guideline: The Claims Administrator did not cite any medical evidence for its decision.

MAXIMUS guideline: Decision based on MTUS Chronic Pain Treatment Guidelines opioids Page(s): 76-84.

Decision rationale: The California chronic pain medical treatment guidelines section on opioids states for ongoing management: On-Going Management. Actions Should Include: (a) Prescriptions from a single practitioner taken as directed, and all prescriptions from a single pharmacy. (b) The lowest possible dose should be prescribed to improve pain and function. (c) Office: Ongoing review and documentation of pain relief, functional status, appropriate medication use, and side effects. Pain assessment should include: current pain; the least reported pain over the period since last assessment; average pain; intensity of pain after taking the opioid; how long it takes for pain relief; and how long pain relief lasts. Satisfactory response to treatment may be indicated by the patient's decreased pain, increased level of function, or improved quality of life. Information from family members or other caregivers should be considered in determining the patient's response to treatment. The 4 A's for Ongoing Monitoring: Four domains have been proposed as most relevant for ongoing monitoring of chronic pain patients on opioids: pain relief, side effects, physical and psychosocial functioning, and the occurrence of any potentially aberrant (or non-adherent) drug-related behaviors. These domains have been summarized as the "4 A's" (analgesia, activities of daily living, adverse side effects, and aberrant drug taking behaviors). The monitoring of these outcomes over time should affect therapeutic decisions and provide a framework for documentation of the clinical use of these controlled drugs. (Passik, 2000) (d) Home: To aid in pain and functioning assessment, the patient should be requested to keep a pain diary that includes entries such as pain triggers, and incidence of end-of-dose pain. It should be emphasized that using this diary will help in tailoring the opioid dose. This should not be a requirement for pain management. (e) Use of drug screening or inpatient treatment with issues of abuse, addiction, or poor pain control. (f) Documentation of misuse of medications (doctor- shopping, uncontrolled drug escalation, drug diversion). (g) Continuing review of overall situation with regard to non-opioid means of pain control. (h) Consideration of a consultation with a multidisciplinary pain clinic if doses of opioids are required beyond what is usually required for the condition or pain does not improve on opioids in 3 months. Consider a psych consult if there is evidence of depression, anxiety or irritability. Consider an addiction medicine consult if there is evidence of substance misuse. When to Continue Opioids (a) If the patient has returned to work. (b) If the patient has improved functioning and pain. (Washington, 2002) (Colorado, 2002) (Ontario, 2000) (VA/DoD, 2003) (Maddox-AAPM/APS, 1997) (Wisconsin, 2004) (Warfield, 2004) The long-term use of this medication class is not recommended per the California MTUS unless there documented evidence of benefit with measurable outcome measures and improvement in function. There is no documented significant decrease in objective pain measures such as VAS scores for significant periods of time. There are no objective measures of improvement of function. Therefore all criteria for the ongoing use of opioids have not been met and the request is not medically necessary.