

| | | | |
|-----------------------|--------------|------------------------------|------------|
| Case Number: | CM15-0112676 | | |
| Date Assigned: | 06/19/2015 | Date of Injury: | 01/19/2015 |
| Decision Date: | 07/17/2015 | UR Denial Date: | 06/01/2015 |
| Priority: | Standard | Application Received: | 06/11/2015 |

HOW THE IMR FINAL DETERMINATION WAS MADE

MAXIMUS Federal Services sent the complete case file to an expert reviewer. He/she has no affiliation with the employer, employee, providers or the claims administrator. He/she has been in active clinical practice for more than five years and is currently working at least 24 hours a week in active practice. The expert reviewer was selected based on his/her clinical experience, education, background, and expertise in the same or similar specialties that evaluate and/or treat the medical condition and disputed items/Service. He/she is familiar with governing laws and regulations, including the strength of evidence hierarchy that applies to Independent Medical Review determinations.

The Expert Reviewer has the following credentials:

State(s) of Licensure: Iowa, Illinois, Hawaii

Certification(s)/Specialty: Preventive Medicine, Occupational Medicine, Public Health & General Preventive Medicine

CLINICAL CASE SUMMARY

The expert reviewer developed the following clinical case summary based on a review of the case file, including all medical records:

The injured worker is a 35 year old female with an industrial injury dated 01/19/2015. The injured worker's diagnoses include pain in joint of shoulder, pain in joint of upper arm, pain in joint of hand, cervicalgia, brachial neuritis or radiculitis not otherwise specified, myalgia and myositis not otherwise specified, sleep disturbance not otherwise specified and skin sensation disturbance. Treatment consisted of diagnostic studies, prescribed medications, and periodic follow up visits. In a progress note dated 05/11/2015, the injured worker reported neck, upper back and right shoulder pain. Cervical exam revealed spasm, tenderness and tight muscle band on the right side. Spinous process tenderness was noted at C5-C7 with tenderness at the paracervical muscles and trapezius. Right shoulder exam revealed restricted range of motion due to pain and tenderness at acromioclavicular joint (AC) and biceps groove. Sensory exam revealed decreased sensation over lateral forearm on the right side. The treating physician prescribed services for Electromyography (EMG) of bilateral upper extremity to determine the origin of her symptoms of radiculopathy.

IMR ISSUES, DECISIONS AND RATIONALES

The Final Determination was based on decisions for the disputed items/services set forth below:

EMG left upper extremity: Upheld

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 177-179.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 11 Forearm, Wrist, and Hand Complaints Page(s): 260-262. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Electrodiagnostic testing (EMG/NCS).

Decision rationale: ACOEM States "Appropriate electrodiagnostic studies (EDS) may help differentiate between CTS and other conditions, such as cervical radiculopathy. These may include nerve conduction studies (NCS), or in more difficult cases, electromyography (EMG) may be helpful." ODG states "Recommended needle EMG or NCS, depending on indications. Surface EMG is not recommended. Electromyography (EMG) and Nerve Conduction Studies (NCS) are generally accepted, well-established and widely used for localizing the source of the neurological symptoms and establishing the diagnosis of focal nerve entrapments, such as carpal tunnel syndrome or radiculopathy, which may contribute to or coexist with CRPS II (causalgia), when testing is performed by appropriately trained neurologists or physical medicine and rehabilitation physicians (improperly performed testing by other providers often gives inconclusive results). As CRPS II occurs after partial injury to a nerve, the diagnosis of the initial nerve injury can be made by electrodiagnostic studies." ODG further clarifies "NCS is not recommended, but EMG is recommended as an option (needle, not surface) to obtain unequivocal evidence of radiculopathy, after 1-month conservative therapy, but EMG's are not necessary if radiculopathy is already clinically obvious." The treating physician does not document evidence of radiculopathy, muscle atrophy, and abnormal neurologic findings. There are no objective findings or subjective complaints for the left upper extremity. The treating physician has not met the above ACOEM and ODG criteria for an EMG of the left upper extremity. As such the request for EMG left upper extremity is not medically necessary.

EMG right upper extremity: Overturned

Claims Administrator guideline: Decision based on MTUS ACOEM Chapter 8 Neck and Upper Back Complaints Page(s): 177-179.

MAXIMUS guideline: Decision based on MTUS ACOEM Chapter 11 Forearm, Wrist, and Hand Complaints Page(s): 260-262. Decision based on Non-MTUS Citation Official Disability Guidelines (ODG) Pain, Electrodiagnostic testing (EMG/NCS).

Decision rationale: ACOEM States "Appropriate electrodiagnostic studies (EDS) may help differentiate between CTS and other conditions, such as cervical radiculopathy. These may include nerve conduction studies (NCS), or in more difficult cases, electromyography (EMG) may be helpful." ODG states "Recommended needle EMG or NCS, depending on indications. Surface EMG is not recommended. Electromyography (EMG) and Nerve Conduction Studies (NCS) are generally accepted, well-established and widely used for localizing the source of the neurological symptoms and establishing the diagnosis of focal nerve entrapments, such as carpal tunnel syndrome or radiculopathy, which may contribute to or coexist with CRPS II (causalgia), when testing is performed by appropriately trained neurologists or physical medicine and

rehabilitation physicians (improperly performed testing by other providers often gives inconclusive results). As CRPS II occurs after partial injury to a nerve, the diagnosis of the initial nerve injury can be made by electrodiagnostic studies." ODG further clarifies "NCS is not recommended, but EMG is recommended as an option (needle, not surface) to obtain unequivocal evidence of radiculopathy, after 1-month conservative therapy, but EMG's are not necessary if radiculopathy is already clinically obvious." The treating physician documents evidence of radiculopathy and abnormal neurologic findings. The patient has also participated in many forms of conservative therapy without significant improvement of subjective complaints. The treating physician has met the above ACOEM and ODG criteria for an EMG of the right upper extremity. As such the request for EMG right upper extremity is medically necessary.